

# Hi95 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP58966

## Product Information

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	<a href="#">P58004</a>
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	54494

## Additional Information

Gene ID	83667
Other Names	Sestrin-2, 1.11.1.-, Hypoxia-induced gene, SESN2 ( <a href="#">HGNC:20746</a> )
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:50-200,Flow-Cyt=2ug/Test,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


## Protein Information

Name	SESN2 ( <a href="#">HGNC:20746</a> )
Function	Functions as an intracellular leucine sensor that negatively regulates the mTORC1 signaling pathway through the GATOR complex (PubMed: <a href="#">18692468</a> , PubMed: <a href="#">25263562</a> , PubMed: <a href="#">25457612</a> , PubMed: <a href="#">26449471</a> , PubMed: <a href="#">26586190</a> , PubMed: <a href="#">26612684</a> , PubMed: <a href="#">31586034</a> , PubMed: <a href="#">35114100</a> , PubMed: <a href="#">35831510</a> , PubMed: <a href="#">36528027</a> ). In absence of leucine, binds the GATOR subcomplex GATOR2 and prevents mTORC1 signaling (PubMed: <a href="#">18692468</a> , PubMed: <a href="#">25263562</a> , PubMed: <a href="#">25457612</a> , PubMed: <a href="#">26449471</a> , PubMed: <a href="#">26586190</a> , PubMed: <a href="#">26612684</a> , PubMed: <a href="#">31586034</a> , PubMed: <a href="#">35114100</a> , PubMed: <a href="#">35831510</a> , PubMed: <a href="#">36528027</a> ). Binding of leucine to SESN2 disrupts its interaction with GATOR2 thereby activating the TORC1 signaling pathway (PubMed: <a href="#">26449471</a> , PubMed: <a href="#">26586190</a> , PubMed: <a href="#">35114100</a> , PubMed: <a href="#">35831510</a> , PubMed: <a href="#">36528027</a> ). This stress-inducible metabolic regulator also plays a role in protection against oxidative and genotoxic stresses. May negatively regulate protein translation in response to endoplasmic reticulum stress, via mTORC1 (PubMed: <a href="#">24947615</a> ). May positively regulate the transcription by

NFE2L2 of genes involved in the response to oxidative stress by facilitating the SQSTM1-mediated autophagic degradation of KEAP1 (PubMed:[23274085](#)). May also mediate TP53 inhibition of TORC1 signaling upon genotoxic stress (PubMed:[18692468](#)). Moreover, may prevent the accumulation of reactive oxygen species (ROS) through the alkylhydroperoxide reductase activity born by the N- terminal domain of the protein (PubMed:[26612684](#)). Was originally reported to contribute to oxidative stress resistance by reducing PRDX1 (PubMed:[15105503](#)). However, this could not be confirmed (PubMed:[19113821](#)).

<b>Cellular Location</b>	Cytoplasm.
<b>Tissue Location</b>	Widely expressed..

## Images



Blank control:K562.

Primary Antibody (green line): Rabbit Anti-Hi95 antibody (AP58966)

Dilution: 2 µg /10<sup>6</sup> cells;


Isotype Control Antibody (orange line): Rabbit IgG .

Secondary Antibody : Goat anti-rabbit IgG-PE

Dilution: 1 µg /test.

Protocol

The cells were fixed with 4% PFA (10min at room temperature)and then permeabilized with 90% ice-cold methanol for 20 min at -20°C. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.



Blank control: K562.

Primary Antibody (green line): Rabbit Anti-Hi95 antibody (AP58966)

Dilution: 2 µg /10<sup>6</sup> cells;

Isotype Control Antibody (orange line): Rabbit IgG .

Secondary Antibody : Goat anti-rabbit IgG-AF488

Dilution: 1 µg /test.

Protocol

The cells were fixed with 4% PFA (10min at room temperature)and then permeabilized with 90% ice-cold methanol for 20 min at -20°C. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.