

CPXM Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP58971

Product Information

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	Q96SM3
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	81668
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human CPXM
Epitope Specificity	251-350/734
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Secreted (By similarity).
SIMILARITY	Belongs to the peptidase M14 family. Contains 1 F5/8 type C domain.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	<p>CPXM (carboxypeptidase X, member 1) belongs to the peptidase M14 family. However, no carboxypeptidase activity has yet been detected. It may be involved in cell-cell interactions. Members of the M14 metalcarboxypeptidase protein family serve many diverse functions and are divided into three subfamilies based on structure, function and amino acid sequence similarity. Belonging to the N/E subfamily, CPXM (metalcarboxypeptidase CPX-1) is a 734 amino acid protein that contains a F5/8 type C domain and likely binds one zinc ion per subunit. Most members of the N/E subfamily contain several domains, including an active carboxypeptidase domain and signal peptide, and are thought to function mostly in protein-protein interactions and/or protein-membrane interactions, thereby targeting the protein to specific locations within the secretory pathway. CPXM is a unique member of this subfamily in that it does not appear to exhibit any enzymatic activity due to lack of several active-site residues that are present in the catalytic domain of other members of the N/E subfamily. Studies showing that CPXM expression is regulated during osteoclastogenesis suggest that CPXM may play a role in osteoclast differentiation. There are two isoforms of CPXM which are a result of alternative splicing events.</p>

Additional Information

Gene ID	56265
Other Names	Probable carboxypeptidase X1, 3.4.17.-, Metallo-carboxypeptidase CPX-1,

CPXM1, CPX1, CPXM

Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:50-200,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	CPXM1
Synonyms	CPX1, CPXM
Function	May be involved in cell-cell interactions. No carboxypeptidase activity was found yet (By similarity).
Cellular Location	Secreted.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.