10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# T2R38 Rabbit pAb

T2R38 Rabbit pAb
Catalog # AP59054

## Product Information

| | |
|---|---|
| **Application** | IHC-P, IHC-F, IF |
| **Primary Accession** | P59533 |
| **Predicted** | Human, Chicken |
| **Host** | Rabbit |
| **Clonality** | Polyclonal |
| **Calculated MW** | 37892 |
| **Physical State** | Liquid |
| **Immunogen** | KLH conjugated synthetic peptide derived from human T2R38/TAS2R38 |
| **Epitope Specificity** | 151-250/333 |
| **Isotype** | IgG |
| **Purity** | affinity purified by Protein A |
| **Buffer** | 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. |
| **SUBCELLULAR LOCATION** | Membrane; Multi-pass membrane protein. |
| **Important Note** | This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications. |
| **Background Descriptions** | The sense of taste is essential for the survival of organisms. For example, the ability to identify sweet-tasting foods enables animals to seek out food with high nutritive value, whereas the ability to identify bitter substances enables them to avoid the ingestion of potentially harmful substances. A family of integral membrane proteins are involved in taste perception and include T1R, which is involved in sweet taste perception and T2R, which is involved in bitter taste perception. Both types of taste receptors couple to various G proteins to initiate signal transduction cascades. Specifically, T2R38 is expressed in subsets of taste receptor cells of the tongue and exclusively in gustducin-positive cells. Variations in T2R38 are associated with the ability to taste the bitter chemical phenylthiocarbamide (PTC), also called thiourea tasting. |

## Additional Information

| | |
|---|---|
| **Gene ID** | 5726 |
| **Other Names** | Taste receptor type 2 member 38, T2R38, PTC bitter taste receptor, Taste receptor type 2 member 61, T2R61, TAS2R38, PTC |
| **Dilution** | IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:5000-10000 |
| **Storage** | Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C. |

## Protein Information

| | |
|---|---|
| **Name** | TAS2R38 |
| **Synonyms** | PTC |
| **Function** | Receptor that may play a role in the perception of bitterness and is gustducin-linked. May play a role in sensing the chemical composition of the gastrointestinal content. The activity of this receptor may stimulate alpha gustducin, mediate PLC-beta-2 activation and lead to the gating of TRPM5 (By similarity). |
| **Cellular Location** | Membrane; Multi-pass membrane protein. |
| **Tissue Location** | Expressed in subsets of taste receptor cells of the tongue and exclusively in gustducin-positive cells. Expressed in testis (PubMed:16720576). |

## Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

## Citations

- Loss of CFTR function is associated with reduced bitter taste receptor-stimulated nitric oxide innate immune responses in nasal epithelial cells and macrophages

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.