

# PISD Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP59091

#### **Product Information**

**Application** WB, IHC-P, IHC-F, IF, ICC, E

Primary Accession
Reactivity
Rat, Pig, Dog
Host
Clonality
Polyclonal
Calculated MW
Physical State

Q9UG56
Rat, Pig, Dog
Rabbit
Polyclonal
Liquid

**Immunogen** KLH conjugated synthetic peptide derived from human PISD

Epitope Specificity 151-250/409

**Isotype** IgG

**Purity** affinity purified by Protein A

**Buffer** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

**SUBCELLULAR LOCATION** Mitochondrion.

**SIMILARITY** Belongs to the phosphatidylserine decarboxylase family.

**SUBUNIT** Heterodimer

**Important Note** This product as supplied is intended for research use only, not for use in

human, therapeutic or diagnostic applications.

**Background Descriptions** Enzymes known as phosphatidylserine decarboxylases (PSDs) catalyze the

formation of phosphatidylethanolamine from phosphatidylserine via phosphatidylserine decarboxylation. Type I PSDs contain LGST motifs and are found in bacteria and eukaryotic mitochondria, whereas type II PSDs contain GGST motifs and are found in eukaryotic endomembrane systems. PISD (phosphatidylserine decarboxylase), also known as phosphatidylserine decarboxylase proenzyme, PSDC, PSD, PSSC, DJ858B16, dJ858B16.2 or DKFZp566G2246, is a 408 amino acid a type I phosphatidylserine

decarboxylase that localizes to the inner mitochondrial membrane. PISD contains a conserved LGST motif which is cleaved to produce two isoforms known as PISD  $\alpha$  and PISD  $\beta$ . PISD is capable of forming a heterodimer and is highly expressed in liver and testis. The gene encoding PISD maps to human

chromosome 22q12.2.

#### **Additional Information**

**Gene ID** 23761

Other Names Phosphatidylserine decarboxylase proenzyme, mitochondrial

{ECO:0000255 | HAMAP-Rule:MF\_03208}, 4.1.1.65

{ECO:0000255|HAMAP-Rule:MF\_03208}, Phosphatidylserine decarboxylase beta chain {ECO:0000255|HAMAP-Rule:MF\_03208}, Phosphatidylserine decarboxylase alpha chain {ECO:0000255|HAMAP-Rule:MF\_03208}, PISD

{ECO:0000255 | HAMAP-Rule:MF\_03208}

**Dilution** WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-50

0,ELISA=1:5000-10000

**Storage** Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

### **Protein Information**

Name PISD {ECO:0000255 | HAMAP-Rule:MF\_03208}

**Function** Catalyzes the formation of phosphatidylethanolamine (PtdEtn) from

phosphatidylserine (PtdSer) (PubMed:30488656, PubMed:30858161). Plays a central role in phospholipid metabolism and in the interorganelle trafficking of phosphatidylserine. May be involved in lipid droplet biogenesis at the

endoplasmic reticulum membrane (By similarity).


**Cellular Location** [Phosphatidylserine decarboxylase beta chain]: Mitochondrion inner

membrane {ECO:0000255 | HAMAP-Rule:MF\_03208,


ECO:0000305 | PubMed:30858161, ECO:0000305 | PubMed:33718843}; Single-pass membrane protein {ECO:0000255 | HAMAP-Rule:MF\_03208}; Intermembrane side {ECO:0000255 | HAMAP-Rule:MF\_03208} [Isoform 1]:

Mitochondrion inner membrane

## **Images**



Paraformaldehyde-fixed, paraffin embedded (Human brain glioma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (PISD) Polyclonal Antibody, Unconjugated (AP59091) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (PISD) Polyclonal Antibody, Unconjugated (AP59091) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.