

Hepcidin-25 Rabbit pAb

Hepcidin-25 Rabbit pAb

Catalog # AP59092

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q99MH3
Predicted	Rat, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	9286
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from rat Hepcidin
Epitope Specificity	60-84/84
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Secreted.
DISEASE	Defects in HAMP are the cause of hemochromatosis type 2B (HFE2B) [MIM:613313]; also known as juvenile hemochromatosis (JH). HFE2B is a disorder of iron metabolism with excess deposition of iron in the tissues, bronze skin pigmentation, hepatic cirrhosis, arthropathy and diabetes. The most common symptoms of hemochromatosis type 2 at presentation are hypogonadism and cardiomyopathy.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The product encoded by this gene is involved in the maintenance of iron homeostasis, and it is necessary for the regulation of iron storage in macrophages, and for intestinal iron absorption. The preproprotein is post-translationally cleaved into mature peptides of 20, 22 and 25 amino acids, and these active peptides are rich in cysteines, which form intramolecular bonds that stabilize their beta-sheet structures. These peptides exhibit antimicrobial activity. Mutations in this gene cause hemochromatosis type 2B, also known as juvenile hemochromatosis, a disease caused by severe iron overload that results in cardiomyopathy, cirrhosis, and endocrine failure. [provided by RefSeq, Jul 2008].

Additional Information

Gene ID	84604
Other Names	Hepcidin, Hamp, Hepc
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:500 0-10000

Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
----------------	---

Protein Information

Name	Hamp
Synonyms	Hepc
Function	Liver-produced hormone that constitutes the main circulating regulator of iron absorption and distribution across tissues. Acts by promoting endocytosis and degradation of ferroportin/SLC40A1, leading to the retention of iron in iron-exporting cells and decreased flow of iron into plasma. Controls the major flows of iron into plasma: absorption of dietary iron in the intestine, recycling of iron by macrophages, which phagocytose old erythrocytes and other cells, and mobilization of stored iron from hepatocytes.
Cellular Location	Secreted.

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.