

# **EF-CBP2 Polyclonal Antibody**

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP59104

## **Product Information**

**Application** WB, IHC-P, IHC-F, IF, E

Primary Accession Q7Z6G3

**Reactivity** Rat, Dog, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 43194

# **Additional Information**

**Gene ID** 54550

Other Names N-terminal EF-hand calcium-binding protein 2, EF-hand calcium-binding

protein 2, Neuronal calcium-binding protein 2, Synaptotagmin-interacting

protein 2, Stip-2, NECAB2, EFCBP2

**Dilution** WB=1:500-2000,IHC-P=1:100-500,IFC-F=1:100-500,IF=1:50-200,ELISA=1:5000-

10000

Format 0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce

**Storage** Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

### **Protein Information**

Name NECAB2

Synonyms EFCBP2

**Function** May act as a signaling scaffold protein that senses intracellular calcium. Can

modulate ligand-induced internalization of ADORA2A and coupling efficiency of mGluR5/GRM5; for both receptors may regulate signaling activity such as

promoting MAPK1/3 (ERK1/2) activation.

Cytoplasm {ECO:0000250 | UniProtKB:F1LQY6,

ECO:0000250 | UniProtKB:Q91ZP9}. Cell projection, dendrite {ECO:0000250 | UniProtKB:F1LQY6}. Cell projection, axon

{ECO:0000250|UniProtKB:F1LQY6}. Cell membrane Note=Colocalizes with

ADORA2A and/or mGluR5/GRM5 at the plasma membrane

(PubMed:17689978, PubMed:19694902). Found in neuronal somata (PubMed:26843217). Detected in the cytoplasm of striatal neurons, at

postsynaptic sites, filling dendritic shafts and spines, and at presynaptic sites, filling axon terminals (By similarity) {ECO:0000250|UniProtKB:F1LQY6, ECO:0000269|PubMed:17689978, ECO:0000269|PubMed:19694902, ECO:0000269|PubMed:26843217}

#### **Tissue Location**

Expressed in brain. Expressed in the spinal dorsal horn with especially strong expression in lamina IIi; found in excitory synaptic boutons and in ependymal cells (at protein level)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.