

PCID1 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP59113

Product Information

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	Q7L2H7
Reactivity	Rat, Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	42503

Additional Information

Gene ID	10480
Other Names	Eukaryotic translation initiation factor 3 subunit M {ECO:0000255 HAMAP-Rule:MF_03012}, eIF3m {ECO:0000255 HAMAP-Rule:MF_03012}, Fetal lung protein B5, hFL-B5, PCI domain-containing protein 1, EIF3M {ECO:0000255 HAMAP-Rule:MF_03012}, HFLB5, PCID1
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:50-200,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	EIF3M {ECO:0000255 HAMAP-Rule:MF_03012}
Synonyms	HFLB5, PCID1
Function	Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis (PubMed: 17403899 , PubMed: 25849773 , PubMed: 27462815). The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl- tRNAi and eIF-5 to form the 43S pre-initiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation (PubMed: 17403899). The eIF-3 complex specifically targets and initiates translation of a subset of mRNAs involved in cell proliferation, including cell

cycling, differentiation and apoptosis, and uses different modes of RNA stem-loop binding to exert either translational activation or repression (PubMed:[25849773](#)).

Cellular Location

Cytoplasm {ECO:0000255|HAMAP-Rule:MF_03012}.

Tissue Location

Broadly expressed..

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.