

RNF17 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP59166

Product Information

Application	IHC-P, IHC-F, IF, E
Primary Accession	Q9BXT8
Reactivity	Rat, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	184643
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human RNF17
Epitope Specificity	1031-1200/1623
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasm (By similarity). Nucleus (By similarity). Note=Predominantly found in the cytoplasm. Component of a nuage in male germ cells (an electron-dense spherical cytoplasmic body present in late pachytene and diplotene spermatocytes and in elonging spermatids) (By similarity).
SIMILARITY	Contains 1 RING-type zinc finger. Contains 4 Tudor domains.
SUBUNIT	Interacts with MXD1, MXD3, MXD4, MXI1 and PIWIL1. Self-associates (By similarity).
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The RING-type zinc finger motif is present in a number of viral and eukaryotic proteins and is made of a conserved cysteine-rich domain that is able to bind two zinc atoms. Proteins that contain this conserved domain are generally involved in the ubiquitination pathway of protein degradation. RNF17 (ring finger protein 17) or tudor domain-containing protein 4, TDRD4, SPATA23, Mmip-2 or FLJ11045, is a testis-specific protein and a novel key regulator of spermiogenesis containing 1,623 amino acids. By distributing Mad proteins to the cytoplasm, RNF17 regulates the transcriptional activity of c-Myc. Although showing localization in the nucleus, RNF17 is predominantly observed in cytoplasm and is a component of a novel nuage found in male germ cells. The gene encoding RNF17 maps to human chromosome 13q12.12 and encodes one RING-type zinc finger and four tudor domains. As a result of alternative splice events, five RNF17 isoforms exist.

Additional Information


Gene ID	56163
Other Names	RING finger protein 17, Tudor domain-containing protein 4, RNF17, TDRD4

Target/Specificity	Testis specific.
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:50-200,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	RNF17
Synonyms	TDRD4
Function	Seems to be involved in regulation of transcriptional activity of MYC. In vitro, inhibits DNA-binding activity of Mad-MAX heterodimers. Can recruit Mad transcriptional repressors (MXD1, MXD3, MXD4 and MXI1) to the cytoplasm. May be involved in spermiogenesis (By similarity).
Cellular Location	Cytoplasm. Nucleus. Note=Predominantly found in the cytoplasm. Component of a nuage in male germ cells (an electron-dense spherical cytoplasmic body present in late pachytene and diplotene spermatocytes and in elongating spermatids) (By similarity).
Tissue Location	Testis specific.

Images



Paraformaldehyde-fixed, paraffin embedded (rat testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (RNF17) Polyclonal Antibody, Unconjugated (AP59166) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.