

RNF138 Rabbit pAb

RNF138 Rabbit pAb

Catalog # AP59197

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q8WVD3
Reactivity	Rat
Predicted	Pig, Dog, Human, Mouse, Rabbit, Horse, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	28193
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human RNF138
Epitope Specificity	51-150/245
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	E3 ubiquitin-protein ligase. Together with NLK, involved in the ubiquitination and degradation of TCF/LEF. Also exhibits auto-ubiquitination activity in combination with UBE2K. May act as a negative regulator in the Wnt/beta-catenin-mediated signaling pathway.

Additional Information

Gene ID	51444
Other Names	E3 ubiquitin-protein ligase RNF138, 2.3.2.27, Nemo-like kinase-associated RING finger protein, NLK-associated RING finger protein, hNARF, RING finger protein 138, RING-type E3 ubiquitin transferase RNF138, RNF138 (HGNC:17765)
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	RNF138 (HGNC:17765)
-------------	---------------------------------------

Function

E3 ubiquitin-protein ligase involved in DNA damage response by promoting DNA resection and homologous recombination (PubMed:[26502055](#), PubMed:[26502057](#)). Recruited to sites of double-strand breaks following DNA damage and specifically promotes double-strand break repair via homologous recombination (PubMed:[26502055](#), PubMed:[26502057](#)). Two different, non-exclusive, mechanisms have been proposed. According to a report, regulates the choice of double-strand break repair by favoring homologous recombination over non-homologous end joining (NHEJ): acts by mediating ubiquitination of XRCC5/Ku80, leading to remove the Ku complex from DNA breaks, thereby promoting homologous recombination (PubMed:[26502055](#)). According to another report, cooperates with UBE2Ds E2 ubiquitin ligases (UBE2D1, UBE2D2, UBE2D3 or UBE2D4) to promote homologous recombination by mediating ubiquitination of RBBP8/CtIP (PubMed:[26502057](#)). Together with NLK, involved in the ubiquitination and degradation of TCF/LEF (PubMed:[16714285](#)). Also exhibits auto-ubiquitination activity in combination with UBE2K (PubMed:[16714285](#)). May act as a negative regulator in the Wnt/beta-catenin-mediated signaling pathway (PubMed:[16714285](#)).


Cellular Location

Chromosome. Note=Recruited at DNA damage sites (PubMed:[26502055](#)). Localizes to sites of double-strand break: localization to double-strand break sites is mediated by the zinc fingers (PubMed:[26502055](#), PubMed:[26502057](#)).

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Paraformaldehyde-fixed, paraffin embedded (rat testis tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (RNF138) Polyclonal Antibody, Unconjugated (AP59197) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.