

# RNF150 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP59198

#### **Product Information**

**Application** WB, IHC-P, IHC-F, IF, ICC, E

Primary Accession

Reactivity

Rost

Clonality

Calculated MW

Physical State

Rabbit

Polyclonal

48072

Liquid

Immunogen KLH conjugated synthetic peptide derived from human RNF150

Epitope Specificity 51-150/347

Isotype IgG

**Purity** affinity purified by Protein A

**Buffer** 0.01M TBS (pH

SUBCELLULAR LOCATION

SIMILARITY Important Note

0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Contains 1 PA (protease associated) domain.Contains 1 RING-type zinc finger.

Contains 1 PA (protease associated) domain. Contains 1 RING-type zinc finger.

Contains 1 PA (protease associated) domain. Contains 1 RING-type zinc finger.

This product as supplied is intended for research use only, not for use in

human, therapeutic or diagnostic applications.

**Background Descriptions** Predicted to enable ubiquitin protein ligase activity. Predicted to be involved

in ubiquitin-dependent protein catabolic process. Predicted to be integral component of membrane. Predicted to be active in cytoplasm. [provided by

Alliance of Genome Resources, Apr 20221

#### **Additional Information**

**Gene ID** 57484

Other Names RING finger protein 150, RNF150, KIAA1214

**Dilution** WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:50-200,

ELISA=1:5000-10000

Format 0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce

**Storage** Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

### **Protein Information**


Name RNF150

Synonyms KIAA1214

Cellular Location

Membrane; Single-pass type I membrane protein

## **Images**




Sample: Brain (Mouse) Lysate at 40 ug

Primary: Anti-RNF150 (AP59198) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at

1/20000 dilution

Predicted band size: 44 kD Observed band size: 44 kD



Sample: Lovo Cell (Human) Lysate at 40 ug Primary: Anti-RNF150 (AP59198) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at

1/20000 dilution

Predicted band size: 44 kD Observed band size: 44 kD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.