

BEND5 Rabbit pAb

BEND5 Rabbit pAb
Catalog # AP59208

Product Information

Primary Accession	Q7L4P6
Reactivity	Mouse
Predicted	Rat, Human, Rabbit, Chicken
Host	Rabbit
Clonality	Polyclonal
Calculated MW	48182
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human BEND5
Epitope Specificity	31-130/421
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Additional Information


Gene ID	79656
Other Names	BEN domain-containing protein 5, BEND5, C1orf165
Dilution	Flow-Cyt=2ug/Test
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	BEND5
Synonyms	C1orf165
Function	Acts as a transcriptional repressor (PubMed: 23468431).

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.



Blank control: Mouse spleen.

Primary Antibody (green line): Rabbit Anti-BEND5 antibody (AP59208)

Dilution: 2 μg / 10^6 cells;


Isotype Control Antibody (orange line): Rabbit IgG .

Secondary Antibody : Goat anti-rabbit IgG-AF488

Dilution: 1 μg /test.

Protocol

The cells were fixed with 70% ethanol (10min at room temperature) and then were incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.



Blank control: Mouse spleen.

Primary Antibody (green line): Rabbit Anti-BEND5 antibody (AP59208)

Dilution: 2 μg / 10^6 cells;

Isotype Control Antibody (orange line): Rabbit IgG .

Secondary Antibody : Goat anti-rabbit IgG-AF488

Dilution: 1 μg /test.

Protocol

The cells were fixed with 70% ethanol (10min at room temperature) and then were incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.