

RNF187 Rabbit pAb

RNF187 Rabbit pAb

Catalog # AP59209

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q5TA31
Predicted	Rat, Human, Mouse, Rabbit
Host	Rabbit
Clonality	Polyclonal
Calculated MW	26190
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human RNF187
Epitope Specificity	101-235/235
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasm. Nucleus. Predominantly located in the cytoplasm. Shuttles between the cytoplasm and the nucleus.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	E3 ubiquitin-protein ligase that acts as a coactivator of JUN-mediated gene activation in response to growth factor signaling via the MAP3K1 pathway, independently from MAPK8.

Additional Information

Gene ID	149603
Other Names	E3 ubiquitin-protein ligase RNF187, 2.3.2.27, RING domain AP1 coactivator 1, RACO-1, RING finger protein 187, RING-type E3 ubiquitin transferase RNF187, RNF187
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	RNF187
Function	E3 ubiquitin-protein ligase that acts as a coactivator of JUN-mediated gene

activation in response to growth factor signaling via the MAP3K1 pathway, independently from MAPK8.

Cellular Location

Cytoplasm. Nucleus. Note=Predominantly located in the cytoplasm. Shuttles between the cytoplasm and the nucleus

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.