

# SPO11 Rabbit pAb

SPO11 Rabbit pAb  
Catalog # AP59210

## Product Information

---

<b>Application</b>	WB
<b>Primary Accession</b>	<a href="#">Q9Y5K1</a>
<b>Reactivity</b>	Human, Mouse, Rat
<b>Predicted</b>	Dog, Pig, Horse
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	44537
<b>Physical State</b>	Liquid
<b>Immunogen</b>	KLH conjugated synthetic peptide derived from human SPO11
<b>Epitope Specificity</b>	151-330/396
<b>Isotype</b>	IgG
<b>Purity</b>	affinity purified by Protein A
<b>Buffer</b>	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
<b>SUBCELLULAR LOCATION</b>	Nuclear
<b>SIMILARITY</b>	Belongs to the TOP6A family.
<b>Important Note</b>	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
<b>Background Descriptions</b>	Spo11 is a type II topoisomerase that is thought to generate the chromosome breaks that initiate meiotic recombination. The Spo11 protein initiates meiotic recombination by generating DNA double-strand breaks (DSBs) and is required for meiotic synapsis in <i>S. cerevisiae</i> . The DSBs are located mostly in promoter regions, where the chromatin is in an open configuration, and cluster in domains along the chromosome. Expression of the Spo11 is detected mainly in the testis, in agreement with its predicted function in the initiation of meiotic recombination. Disruption of Spo11 leads to severe gonadal abnormalities from defective meiosis and results in infertility.

## Additional Information

---

<b>Gene ID</b>	23626
<b>Other Names</b>	Meiotic recombination protein SPO11, 5.6.2.2, Cancer/testis antigen 35, CT35, SPO11
<b>Target/Specificity</b>	Highly expressed in testis.
<b>Dilution</b>	WB=1:500-2000
<b>Storage</b>	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


## Protein Information

<b>Name</b>	SPO11
<b>Function</b>	Component of a topoisomerase 6 complex specifically required for meiotic recombination. Together with TOP6BL, mediates DNA cleavage that forms the double-strand breaks (DSB) that initiate meiotic recombination. The complex promotes relaxation of negative and positive supercoiled DNA and DNA decatenation through cleavage and ligation cycles. Essential for the phosphorylation of SMC3, HORMAD1 and HORMAD2.
<b>Cellular Location</b>	Nucleus.
<b>Tissue Location</b>	Highly expressed in testis.

## Background

Spo11 is a type II topoisomerase that is thought to generate the chromosome breaks that initiate meiotic recombination. The Spo11 protein initiates meiotic recombination by generating DNA double-strand breaks (DSBs) and is required for meiotic synapsis in *S. cerevisiae*. The DSBs are located mostly in promoter regions, where the chromatin is in an open configuration, and cluster in domains along the chromosome. Expression of the Spo11 is detected mainly in the testis, in agreement with its predicted function in the initiation of meiotic recombination. Disruption of Spo11 leads to severe gonadal abnormalities from defective meiosis and results in infertility.

## Images



**Sample:**  
Lane 1: Mouse Testis tissue lysates  
Lane 2: Rat Testis tissue lysates  
Lane 3: Human PC-3 cell lysates  
Primary: Anti-SPO11 (AP59210) at 1/1000 dilution  
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution  
Predicted band size: 45 kDa  
Observed band size: 48 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.