

# UFD1L Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP59366

## Product Information

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	<a href="#">Q92890</a>
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	34500

## Additional Information

Gene ID	7353
Other Names	Ubiquitin recognition factor in ER-associated degradation protein 1 {ECO:0000312 HGNC:HGNC:12520}, Ubiquitin fusion degradation protein 1, UB fusion protein 1, UFD1 ( <a href="#">HGNC:12520</a> ), UFD1L
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:50-200,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

## Protein Information

Name	UFD1 ( <a href="#">HGNC:12520</a> )
Synonyms	UFD1L
Function	Essential component of the ubiquitin-dependent proteolytic pathway which degrades ubiquitin fusion proteins. The ternary complex containing UFD1, VCP and NPLOC4 binds ubiquitinated proteins and is necessary for the export of misfolded proteins from the ER to the cytoplasm, where they are degraded by the proteasome. The NPLOC4-UFD1- VCP complex regulates spindle disassembly at the end of mitosis and is necessary for the formation of a closed nuclear envelope. It may be involved in the development of some ectoderm-derived structures (By similarity). Acts as a negative regulator of type I interferon production via the complex formed with VCP and NPLOC4, which binds to RIGI and recruits RNF125 to promote ubiquitination and degradation of RIGI (PubMed: <a href="#">26471729</a> ).

<b>Cellular Location</b>	Nucleus {ECO:0000250 UniProtKB:Q9ES53}. Cytoplasm, cytosol {ECO:0000250 UniProtKB:Q9ES53}
<b>Tissue Location</b>	Found in adult heart, skeletal muscle and pancreas, and in fetal liver and kidney

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.