10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

CNTD Rabbit pAb

CNTD Rabbit pAb
Catalog # AP59368

Product Information

Application	WB
Primary Accession	Q8N815
Reactivity	Mouse
Predicted	Rat, Pig, Human, Horse, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	36921
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human CNTD
Epitope Specificity	1-100/330
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	No data available.

Additional Information

Gene ID	124817
Other Names	Cyclin N-terminal domain-containing protein 1, CNTD1 (HGNC:26847), CNTD
Dilution	WB=1:500-2000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information


Name	CNTD1 (HGNC:26847)
Synonyms	CNTD
Function	Plays a role in the different steps of crossover formation during meiotic recombination. Participates in the crossover differentiation step of crossover-specific recombination intermediates through its interaction with PRR19. In addition, stimulates crossover formation through the interactions

	with RFC3 and RFC4 and simultaneously regulates cell-cycle progression through interactions with CDC34 and subsequent ubiquitination of WEE1. May also participates in an active deselection process that destabilizes or removes excess pre-CO intermediates.
Cellular Location	Nucleus {ECO:0000250|UniProtKB:Q9D995}. Cytoplasm {ECO:0000250|UniProtKB:Q9D995}. Chromosome {ECO:0000250|UniProtKB:Q9D995}. Note=Shuttles between the nucleus and cytoplasm in a stage-specific manner of prophase I cells. Co-localized at crossover sites with PRR19. {ECO:0000250|UniProtKB:Q9D995}

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Sample: Brain (Mouse) Lysate at 40 ug
Primary: Anti-CNTD (AP59368) at 1/300 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 37 kD
Observed band size: 34 kD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.