

## C10orf93 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP59379

## **Product Information**

**Application** WB, IHC-P, IHC-F, IF, E

Primary Accession

Reactivity
Rat, Bovine
Host
Rabbit
Clonality
Polyclonal
Calculated MW
303500
Physical State
Liquid

**Immunogen** KLH conjugated synthetic peptide derived from human C10orf93

Epitope Specificity 951-1050/2715

**Isotype** IgG

**Purity** affinity purified by Protein A

**Buffer** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

**SIMILARITY** Contains 13 TPR repeats.

**Important Note** This product as supplied is intended for research use only, not for use in

human, therapeutic or diagnostic applications.

**Background Descriptions** C10orf93 , also known as C10orf124 or TPR repeat-containing protein

C10orf93, is a 1,530 amino acid protein that contains two TPR repeats and exists as three alternatively spliced isoforms. The gene encoding C10orf93 maps to human chromosome 10q26.3. Spanning nearly 135 million base pairs, chromosome 10 makes up approximately 4.5% of total DNA in cells and encodes nearly 1,200 genes. Several protein-coding genes, including those that encode for chemokines, cadherins, excision repair proteins, early growth response factors (Egrs) and fibroblast growth receptors (FGFRs), are located on chromosome 10. Defects in some of the genes that map to chromosome 10 are associated with Charcot-Marie Tooth disease, Jackson-Weiss syndrome, Usher syndrome, nonsyndromatic deafness, Wolman's syndrome, Cowden

syndrome, multiple endocrine neoplasia type 2 and porphyria.

## **Additional Information**

**Gene ID** 54777

Other Names Cilia- and flagella-associated protein 46, Tetratricopeptide repeat protein 40

{ECO:0000312 | HGNC:HGNC:25247}, CFAP46 (HGNC:25247)

**Dilution** WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:50-200,ELISA=1:5000-

10000

Format 0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce

**Storage** Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4  $^{\circ}$ C.

## **Protein Information**

Name CFAP46 ( HGNC:25247)

**Function** As part of the central apparatus of the cilium axoneme plays a role in cilium

movement.

**Cellular Location** Cytoplasm, cytoskeleton, cilium axoneme {ECO:0000250 | UniProtKB:A8ICS9}

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.