

PDCL3 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP59446

Product Information

Application	IHC-P, IHC-F, IF, ICC
Primary Accession	Q9H2J4
Reactivity	Rat, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	27614

Additional Information

Gene ID	79031
Other Names	Phosducin-like protein 3, HTPHP, PhPL3, Viral IAP-associated factor 1, VIAF-1, PDCL3, PhLP2A, VIAF1
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	PDCL3
Synonyms	PhLP2A, VIAF1
Function	Acts as a chaperone for the angiogenic VEGF receptor KDR/VEGFR2, increasing its abundance by inhibiting its ubiquitination and degradation (PubMed: 23792958 , PubMed: 26059764). Inhibits the folding activity of the chaperonin-containing T-complex (CCT) which leads to inhibition of cytoskeletal actin folding (PubMed: 17429077). Acts as a chaperone during heat shock alongside HSP90 and HSP40/70 chaperone complexes (By similarity). Modulates the activation of caspases during apoptosis (PubMed: 15371430).
Cellular Location	Cytoplasm. Cytoplasm, perinuclear region. Endoplasmic reticulum
Tissue Location	Expressed in endothelial cells (at protein level) (PubMed:26059764). Expressed in all tissues examined including spleen, thymus, prostate, testis, ovary, small intestine and colon (PubMed:15371430).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.