

Anti-ARF6 Antibody

Rabbit polyclonal antibody to ARF6

Catalog # AP59481

Product Information

Application	WB
Primary Accession	P62330
Other Accession	P62331
Reactivity	Human, Mouse, Rat, Zebrafish, Pig, Chicken
Host	Rabbit
Clonality	Polyclonal
Calculated MW	20082

Additional Information

Gene ID	382
Other Names	ADP-ribosylation factor 6
Target/Specificity	Recognizes endogenous levels of ARF6 protein.
Dilution	WB~~WB (1/500 - 1/1000)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	ARF6 {ECO:0000303 Ref.6, ECO:0000312 HGNC:HGNC:659}
Function	GTP-binding protein involved in protein trafficking that regulates endocytic recycling and cytoskeleton remodeling (PubMed: 11266366 , PubMed: 16737952 , PubMed: 18400762 , PubMed: 21170023 , PubMed: 32103017 , PubMed: 7589240). GTP-bound form plays an important role in the transport of multiple palmitoylated proteins from the Golgi to the plasma membrane (PubMed: 37461827). Required for normal completion of mitotic cytokinesis (By similarity). Plays a role in the reorganization of the actin cytoskeleton and the formation of stress fibers (By similarity). Involved in the regulation of dendritic spine development, contributing to the regulation of dendritic branching and filopodia extension (PubMed: 14978216). Potentiates the neurite outgrowth in primary neurons by interacting with the molecular adapter APBB1 (PubMed: 36250347). Plays an important role in membrane trafficking, during junctional remodeling and epithelial polarization (PubMed: 36017701). Regulates surface levels of adherens junction proteins such as CDH1 (By similarity). Required for NTRK1

sorting to the recycling pathway from early endosomes (By similarity).

Cellular Location

Cytoplasm, cytosol. Cell membrane; Lipid-anchor. Endosome membrane; Lipid-anchor. Recycling endosome membrane; Lipid-anchor. Cell projection, filopodium membrane; Lipid- anchor. Cell projection, ruffle. Cleavage furrow. Midbody, Midbody ring. Early endosome membrane {ECO:0000250|UniProtKB:P62331}; Lipid-anchor {ECO:0000250|UniProtKB:P62331}. Golgi apparatus, trans-Golgi network membrane {ECO:0000250|UniProtKB:P62331}; Lipid-anchor {ECO:0000250|UniProtKB:P62331}. Note=Distributed uniformly on the plasma membrane, as well as throughout the cytoplasm during metaphase Subsequently concentrated at patches in the equatorial region at the onset of cytokinesis, and becomes distributed in the equatorial region concurrent with cleavage furrow ingression. In late stages of cytokinesis, concentrates at the midbody ring/Flemming body (PubMed:23603394). Recruitment to the midbody ring requires both activation by PSD/EFA6A and interaction with KIF23/MKLP1 (PubMed:23603394). After abscission of the intercellular bridge, incorporated into one of the daughter cells as a midbody remnant and localizes to punctate structures beneath the plasma membrane (PubMed:23603394). Recruited to the cell membrane in association with CYTH2 and ARL4C (PubMed:17398095). Colocalizes with DAB2IP at the plasma membrane and endocytic vesicles (PubMed:19948740) Myristoylation is required for proper localization to membranes: myristoylation on Lys-3 allows ARF6 to remain on membranes during the GTPase cycle (PubMed:32103017, PubMed:7589240)


Tissue Location

Ubiquitous, with higher levels in heart, substantia nigra, and kidney.

Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human ARF6. The exact sequence is proprietary.

Images



Western blot analysis of ARF6 expression in HEK293T (A), Hela (B), A2780 (C), mouse liver (D), mouse spleen (E), rat liver (F), rat spleen (G) whole cell lysates.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.