

# Anti-CstF-64 Antibody

Rabbit polyclonal antibody to CstF-64 Catalog # AP59527

#### **Product Information**

**Application** WB, IP, IF/IC, IHC

Primary Accession P33240
Other Accession O8BIO5

**Reactivity** Human, Mouse, Rat, Zebrafish, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 60959

#### **Additional Information**

Gene ID 1478

Other Names Cleavage stimulation factor subunit 2; CF-1 64 kDa subunit; Cleavage

stimulation factor 64 kDa subunit; CSTF 64 kDa subunit; CstF-64

**Target/Specificity** KLH-conjugated synthetic peptide encompassing a sequence within the

N-term region of human CstF-64. The exact sequence is proprietary.

**Dilution** WB~~WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500), IP (1/10 -

1/100) IP~~N/A IF/IC~~N/A IHC~~WB (1/500 - 1/1000), IHC (1/100 - 1/200),

IF/IC (1/100 - 1/500), IP (1/10 - 1/100)

**Format** Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30%

glycerol, and 0.09% (W/V) sodium azide.

**Storage** Store at -20 °C.Stable for 12 months from date of receipt

#### **Protein Information**

Name CSTF2

**Function** One of the multiple factors required for polyadenylation and 3'-end cleavage

of mammalian pre-mRNAs. This subunit is directly involved in the binding to


pre-mRNAs.

**Cellular Location** Nucleus. Note=Localized with DDX1 in cleavage bodies.


## **Background**

KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human CstF-64.


### **Images**



Western blot analysis of CstF-64 expression in mouse spleen (A), rat spleen (B), rat brain (C) whole cell lysates.



Immunohistochemical analysis of CstF-64 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of CstF-64 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.