

Anti-DGK alpha Antibody

Rabbit polyclonal antibody to DGK alpha

Catalog # AP59533

Product Information

Application	WB, IF/IC, IHC
Primary Accession	P23743
Reactivity	Human, Monkey
Host	Rabbit
Clonality	Polyclonal
Calculated MW	82630

Additional Information

Gene ID	1606
Other Names	DAGK; DAGK1; Diacylglycerol kinase alpha; DAG kinase alpha; 80 kDa diacylglycerol kinase; Diglyceride kinase alpha; DGK-alpha
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human DGK alpha. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500) IF/IC~~N/A IHC~~WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	DGKA
Synonyms	DAGK, DAGK1
Function	Diacylglycerol kinase that converts diacylglycerol/DAG into phosphatidic acid/phosphatidate/PA and regulates the respective levels of these two bioactive lipids (PubMed: 15544348 , PubMed: 2175712). Thereby, acts as a central switch between the signaling pathways activated by these second messengers with different cellular targets and opposite effects in numerous biological processes (PubMed: 15544348 , PubMed: 2175712). Also plays an important role in the biosynthesis of complex lipids (Probable). Can also phosphorylate 1-alkyl-2- acylglycerol in vitro as efficiently as diacylglycerol provided it contains an arachidonoyl group (PubMed: 15544348). Also involved in the production of alkyl-lysophosphatidic acid, another bioactive lipid,

through the phosphorylation of 1-alkyl-2-acetyl glycerol (PubMed:[22627129](#)).

Cellular Location

Cytoplasm, cytosol.


Tissue Location

Expressed in lymphocytes.


Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human DGK alpha. The exact sequence is proprietary.


Images



Western blot analysis of DGK alpha expression in EC9706 (A), Myla2059 (B) whole cell lysates.



Immunohistochemical analysis of DGK alpha staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of DGK alpha staining in MDCK cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.