

# Anti-GRK5 Antibody

Rabbit polyclonal antibody to GRK5  
Catalog # AP59568

## Product Information

---

Application	WB, IP, IHC
Primary Accession	<a href="#">P34947</a>
Other Accession	<a href="#">Q8VEB1</a>
Reactivity	Human, Mouse, Rat, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	67787

## Additional Information

---

Gene ID	2869
Other Names	GPRK5; G protein-coupled receptor kinase 5; G protein-coupled receptor kinase GRK5
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human GRK5. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000), IHC (1/100 - 1/200), IP (1/10 - 1/100) IP~~N/A IHC~~WB (1/500 - 1/1000), IHC (1/100 - 1/200), IP (1/10 - 1/100)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

## Protein Information

---

Name	GRK5
Synonyms	GPRK5
Function	Serine/threonine kinase that phosphorylates preferentially the activated forms of a variety of G-protein-coupled receptors (GPCRs). Such receptor phosphorylation initiates beta-arrestin-mediated receptor desensitization, internalization, and signaling events leading to their down-regulation. Phosphorylates a variety of GPCRs, including adrenergic receptors, muscarinic acetylcholine receptors (more specifically Gi-coupled M2/M4 subtypes), dopamine receptors and opioid receptors. In addition to GPCRs, also phosphorylates various substrates: Hsc70-interacting protein/ST13, TP53/p53, HDAC5, and arrestin-1/ARRB1. Phosphorylation of ARRB1 by GRK5 inhibits G-protein independent MAPK1/MAPK3 signaling downstream of

5HT4-receptors. Phosphorylation of HDAC5, a repressor of myocyte enhancer factor 2 (MEF2) leading to nuclear export of HDAC5 and allowing MEF2-mediated transcription. Phosphorylation of TP53/p53, a crucial tumor suppressor, inhibits TP53/p53-mediated apoptosis. Phosphorylation of ST13 regulates internalization of the chemokine receptor. Phosphorylates rhodopsin (RHO) (in vitro) and a non G-protein-coupled receptor, LRP6 during Wnt signaling (in vitro).

#### Cellular Location

Cytoplasm. Nucleus. Cell membrane; Peripheral membrane protein.  
Note=Predominantly localized at the plasma membrane; targeted to the cell surface through the interaction with phospholipids. Nucleus localization is regulated in a GPCR and Ca(2+)/calmodulin-dependent fashion


#### Tissue Location

Highest levels in heart, placenta, lung > skeletal muscle > brain, liver, pancreas > kidney.


## Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human GRK5. The exact sequence is proprietary.

## Images



Western blot analysis of GRK5 expression in Hela (A), H446 (B), mouse kidney (C), mouse lung (D), rat kidney (E) whole cell lysates.



Immunohistochemical analysis of GRK5 staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.