

# Anti-GLUR2 Antibody

Rabbit polyclonal antibody to GLUR2 Catalog # AP59569

#### **Product Information**

ApplicationWBPrimary AccessionP42262Other AccessionP23819

**Reactivity** Human, Mouse, Rat, Zebrafish, Monkey, Chicken

HostRabbitClonalityPolyclonalCalculated MW98821

### **Additional Information**

**Gene ID** 2891

Other Names GLUR2; Glutamate receptor 2; GluR-2; AMPA-selective glutamate receptor 2;

GluR-B; GluR-K2; Glutamate receptor ionotropic, AMPA 2; GluA2

**Target/Specificity** Recognizes endogenous levels of GLUR2 protein.

**Dilution** WB~~WB (1/500 - 1/1000)

**Format** Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30%

glycerol, and 0.09% (W/V) sodium azide.

Storage Store at -20 °C.Stable for 12 months from date of receipt

#### **Protein Information**

Name GRIA2 (<u>HGNC:4572</u>)

**Function** Ionotropic glutamate receptor that functions as a ligand- gated cation

channel, gated by L-glutamate and glutamatergic agonists such as

alpha-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA), quisqualic

acid, and kainic acid (PubMed:20614889, PubMed:31300657,

PubMed:8003671). L-glutamate acts as an excitatory neurotransmitter at many synapses in the central nervous system and plays an important role in fast excitatory synaptic transmission (PubMed:14687553). Binding of the excitatory neurotransmitter L- glutamate induces a conformation change, leading to the opening of the cation channel, and thereby converts the chemical signal to an electrical impulse upon entry of monovalent and divalent cations such as sodium and calcium (PubMed:20614889,

PubMed:<u>8003671</u>). The receptor then desensitizes rapidly and enters in a transient inactive state, characterized by the presence of bound agonist (By similarity). In the presence of CACNG4 or CACNG7 or CACNG8, shows

resensitization which is characterized by a delayed accumulation of current flux upon continued application of L-glutamate (By similarity). Through complex formation with NSG1, GRIP1 and STX12 controls the intracellular fate of AMPAR and the endosomal sorting of the GRIA2 subunit toward recycling and membrane targeting (By similarity).


#### **Cellular Location**

Cell membrane; Multi-pass membrane protein. Postsynaptic cell membrane; Multi-pass membrane protein. Postsynaptic density membrane {ECO:0000250 | UniProtKB:P23819}; Multi-pass membrane protein {ECO:0000250 | UniProtKB:P23819}. Note=Interaction with CACNG2, CNIH2 and CNIH3 promotes cell surface expression (By similarity). Displays a somatodendritic localization and is excluded from axons in neurons (By similarity). {ECO:0000250 | UniProtKB:P19491, ECO:0000250 | UniProtKB:P23819}

# **Background**

KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human GLUR2. The exact sequence is proprietary.

## **Images**



Western blot analysis of GLUR2 expression in mouse brain (A), rat brain (B) whole cell lysates.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.