

Anti-c-Jun (pY170) Antibody

Rabbit polyclonal antibody to c-Jun (pY170) Catalog # AP59600

Product Information

Application WB, IP, IHC
Primary Accession P05412
Other Accession P05627

Reactivity Human, Mouse, Rat, Pig, Chicken, Bovine

HostRabbitClonalityPolyclonalCalculated MW35676

Additional Information

Gene ID 3725

Other Names Transcription factor AP-1; Activator protein 1; AP1; Proto-oncogene c-Jun;

V-jun avian sarcoma virus 17 oncogene homolog; p39

Target/Specificity KLH-conjugated synthetic peptide encompassing a sequence within the center

region of human c-Jun. The exact sequence is proprietary.

Dilution WB~~WB (1/500 - 1/1000), IHC (1/100 - 1/200), IP (1/10 - 1/100) IP~~N/A

IHC~~WB (1/500 - 1/1000), IHC (1/100 - 1/200), IP (1/10 - 1/100)

Format Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30%

glycerol, and 0.09% (W/V) sodium azide.

Storage Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name JUN

Function Transcription factor that recognizes and binds to the AP-1 consensus motif

5'-TGA[GC]TCA-3' (PubMed:10995748, PubMed:22083952). Heterodimerizes with proteins of the FOS family to form an AP-1 transcription complex, thereby enhancing its DNA binding activity to the AP-1 consensus sequence 5'-TGA[GC]TCA-3' and enhancing its transcriptional activity (By similarity). Together with FOSB, plays a role in activation-induced cell death of T cells by binding to the AP-1 promoter site of FASLG/CD95L, and inducing its transcription in response to activation of the TCR/CD3 signaling pathway (PubMed:12618758). Promotes activity of NR5A1 when phosphorylated by HIPK3 leading to increased steroidogenic gene expression upon cAMP signaling pathway stimulation (PubMed:17210646). Involved in activated

KRAS-mediated transcriptional activation of USP28 in colorectal cancer (CRC)

cells (PubMed:<u>24623306</u>). Binds to the USP28 promoter in colorectal cancer (CRC) cells (PubMed:<u>24623306</u>).

Cellular Location

Nucleus.


Tissue Location

Expressed in the developing and adult prostate and prostate cancer cells.


Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human c-Jun. The exact sequence is proprietary.

Images



Western blot analysis of c-Jun (pY170) expression in Hela (A), mouse kidney (B) whole cell lysates.



Immunohistochemical analysis of c-Jun (pY170) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.