

Anti-OCT3 Antibody

Rabbit polyclonal antibody to 41915 Catalog # AP59702

Product Information

Application WB, IP **Primary Accession** 075751

Reactivity Human, Mouse, Rat

HostRabbitClonalityPolyclonalCalculated MW61280

Additional Information

Gene ID 6581

Other Names EMTH; OCT3; Solute carrier family 22 member 3; Extraneuronal monoamine

transporter; EMT; Organic cation transporter 3

Target/Specificity Recognizes endogenous levels of OCT3 protein.

Dilution WB~~WB (1/500 - 1/1000), IP (1/10 - 1/100) IP~~N/A

Format Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30%

glycerol, and 0.09% (W/V) sodium azide.

Storage Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name SLC22A3 (<u>HGNC:10967</u>)

Function Electrogenic voltage-dependent transporter that mediates the transport of a

variety of organic cations such as endogenous bioactive amines, cationic

drugs and xenobiotics (PubMed: 10196521, PubMed: 10966924,

PubMed: 12538837, PubMed: 17460754, PubMed: 20858707). Cation cellular uptake or release is driven by the electrochemical potential, i.e. membrane potential and concentration gradient (PubMed: 10966924). Functions as a Na(+)- and Cl(-)-independent, bidirectional uniporter (PubMed: 12538837). Implicated in monoamine neurotransmitters uptake such as dopamine, adrenaline/epinephrine, noradrenaline/norepinephrine, histamine, serotonin and tyramine, thereby supporting a role in homeostatic regulation of

aminergic neurotransmission in the brain (PubMed:10196521, PubMed:16581093, PubMed:20858707). Transports dopaminergic neuromodulators cyclo(his- pro) and salsolinol with low efficiency

(PubMed:<u>17460754</u>). May be involved in the uptake and disposition of cationic compounds by renal clearance from the blood flow (PubMed:<u>10966924</u>). May

contribute to regulate the transport of cationic compounds in testis across the blood-testis-barrier (Probable). Mediates the transport of polyamine spermidine and putrescine (By similarity). Mediates the bidirectional transport of polyamine agmatine (PubMed:12538837). Also transports guanidine (PubMed:10966924). May also mediate intracellular transport of organic cations, thereby playing a role in amine metabolism and intracellular signaling (By similarity).

Cellular Location

Cell membrane; Multi-pass membrane protein. Apical cell membrane; Multi-pass membrane protein. Basolateral cell membrane; Multi-pass membrane protein. Mitochondrion membrane {ECO:0000250|UniProtKB:O88446}. Endomembrane system {ECO:0000250|UniProtKB:O88446}. Nucleus membrane {ECO:0000250|UniProtKB:O88446}. Nucleus outer membrane {ECO:0000250|UniProtKB:O88446}. Note=Localized to the apical/brush border membrane of enterocytes (PubMed:16263091). Localized to the luminal/apical membrane of ciliated epithelial cells in bronchi (PubMed:15817714). Localized to the basolateral membrane of intermediate cells in bronchi (PubMed:15817714). Localized to the entire plasma membrane of basal cells in bronchi (PubMed:15817714)


Tissue Location

Expressed in liver (PubMed:10196521, PubMed:9933568). Expressed in intestine (PubMed:16263091, PubMed:20858707). Expressed in kidney in proximal tubular cells (PubMed:10966924). Expressed in placenta (PubMed:10966924, PubMed:9933568). Expressed throughout the brain, including cerebral cortex, cerebrellum, substancia nigra, medulla oblongata, hippocampus, caudate nucleus, nucleus accumbens and pons with low levels of expression detected in nearly all brain regions (PubMed:10196521, PubMed:20858707). In testis, mostly localized to peritubular myoid cells and Leydig cells, and weakly expressed in developing germ cells (PubMed:35307651). Expressed in tracheal and bronchial epithelium of the respiratory tract, where it localizes to the apical membrane of ciliated cells, the entire membrane of basal cells and the basolateral membrane of intermediate cells (PubMed:15817714). Expressed in skeletal muscle, adrenal gland, heart, prostate, aorta, salivary gland, adrenal gland, uterus, lymph node, lung, trachea and spinal cord (PubMed:10196521, PubMed:20858707, PubMed:9933568). Expressed in fetal lung and liver (PubMed:9933568).

Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human 41915. The exact sequence is proprietary.

Images



Western blot analysis of OCT3 expression in HEK293T (A), Hela (B), DLD (C), mouse muscle (D), mouse liver (E), rat muscle (F) whole cell lysates.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.