

Anti-MADD Antibody

Catalog # AP59748

Product Information

Application	WB, IHC, IF/IC
Primary Accession	Q8WVG6
Reactivity	Human, Rat, Mk
Host	Rabbit
Clonality	Polyclonal
Calculated MW	183303

Additional Information

Gene ID	8567
Other Names	DENN; IG20; KIAA0358; MAP kinase-activating death domain protein; Differentially expressed in normal and neoplastic cells; Insulinoma glucagonoma clone 20; Rab3 GDP/GTP exchange factor
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human MADD. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000), IP (1/10 - 1/100) IHC~~1:100~500 IF/IC~~N/A
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	MADD {ECO:0000312 EMBL:AAB57735.1, ECO:0000312 HGNC:HGNC:6766}
Function	Guanyl-nucleotide exchange factor that regulates small GTPases of the Rab family (PubMed: 18559336 , PubMed: 20937701). Converts GDP-bound inactive form of RAB27A and RAB27B to the GTP-bound active forms (PubMed: 18559336 , PubMed: 20937701). Converts GDP-bound inactive form of RAB3A, RAB3C and RAB3D to the GTP-bound active forms, GTPases involved in synaptic vesicle exocytosis and vesicle secretion (By similarity). Plays a role in synaptic vesicle formation and in vesicle trafficking at the neuromuscular junction (By similarity). Involved in up-regulating a post-docking step of synaptic exocytosis in central synapses (By similarity). Probably by binding to the motor proteins KIF1B and KIF1A, mediates motor-dependent transport of GTP-RAB3A- positive vesicles to the presynaptic nerve terminals (By similarity). Plays a role in TNFA-mediated activation of the MAPK pathway, including ERK1/2 (PubMed: 32761064). May link TNFRSF1A with MAP kinase activation (PubMed: 9115275). May be

involved in the regulation of TNFA-induced apoptosis (PubMed:[11577081](#), PubMed:[32761064](#)).

Cellular Location

Cell membrane. Cytoplasm. Cell projection, axon
{ECO:0000250|UniProtKB:Q80U28}


Tissue Location

Expressed in testis, ovary, brain and heart (PubMed:8988362). Expressed in spleen, thymus, prostate, testis, ovary, small intestine and colon (PubMed:9115275). Expressed in liver (PubMed:9796103). [Isoform 2]: Expressed in the brain, breast, kidney, lung, ovary, pancreas, testis, uterus, stomach and thyroid [Isoform 4]: Expressed in the brain, breast, kidney, lung, ovary, pancreas, testis, uterus, stomach and thyroid [Isoform 6]: Not detected in the brain, breast, kidney, lung, ovary, pancreas, testis, uterus, stomach and thyroid


Images



Western blot analysis of MADD expression in PC12 (A) whole cell lysates. (Predicted band size: 183 kD; Observed band size: 200 kD)



Immunohistochemical analysis of MADD staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of MADD staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a AF594-conjugated secondary antibody (red) in PBS at room temperature in the dark.