

# Anti-Neuropilin 1 Antibody

Rabbit polyclonal antibody to Neuropilin 1  
Catalog # AP59758

## Product Information

---

Application	WB
Primary Accession	<a href="#">O14786</a>
Reactivity	Human, Rat, Monkey
Host	Rabbit
Clonality	Polyclonal
Calculated MW	103134

## Additional Information

---

Gene ID	8829
Other Names	NRP; VEGF165R; Neuropilin-1; Vascular endothelial cell growth factor 165 receptor; CD304
Target/Specificity	Recognizes endogenous levels of Neuropilin 1 protein.
Dilution	WB~~WB (1/500 - 1/1000)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

## Protein Information

---

Name	NRP1 ( <a href="#">HGNC:8004</a> )
Synonyms	NRP, VEGF165R
Function	Cell-surface receptor involved in the development of the cardiovascular system, in angiogenesis, in the formation of certain neuronal circuits and in organogenesis outside the nervous system. Mediates the chemorepulsant activity of semaphorins (PubMed: <a href="#">10688880</a> , PubMed: <a href="#">9288753</a> , PubMed: <a href="#">9529250</a> ). Recognizes a C-end rule (CendR) motif R/KXXR/K on its ligands which causes cellular internalization and vascular leakage (PubMed: <a href="#">19805273</a> ). It binds to semaphorin 3A, the PLGF-2 isoform of PGF, the VEGF165 isoform of VEGFA and VEGFB (PubMed: <a href="#">10688880</a> , PubMed: <a href="#">19805273</a> , PubMed: <a href="#">9288753</a> , PubMed: <a href="#">9529250</a> ). Coexpression with KDR results in increased VEGF165 binding to KDR as well as increased chemotaxis. Regulates VEGF-induced angiogenesis. Binding to VEGFA initiates a signaling pathway needed for motor neuron axon guidance and cell body migration, including for the caudal migration of facial motor neurons from

rhombomere 4 to rhombomere 6 during embryonic development (By similarity). Regulates mitochondrial iron transport via interaction with ABCB8/MITOSUR (PubMed:[30623799](#)).

**Cellular Location**

[Isoform 2]: Secreted

**Tissue Location**

[Isoform 1]: The expression of isoforms 1 and 2 does not seem to overlap. Expressed in olfactory epithelium (at protein level) (PubMed:33082293). Expressed in fibroblasts (at protein level) (PubMed:36213313). Expressed by the blood vessels of different tissues In the developing embryo it is found predominantly in the nervous system. In adult tissues, it is highly expressed in heart and placenta; moderately in lung, liver, skeletal muscle, kidney and pancreas; and low in adult brain (PubMed:10688880, PubMed:9529250). Expressed in the central nervous system, including olfactory related regions such as the olfactory tubercles and paraolfactory gyri (PubMed:33082293)


## Background

---

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Neuropilin 1. The exact sequence is proprietary.

## Images

---



Western blot analysis of Neuropilin 1 expression in mouse brain (A), PC3 (B) whole cell lysates.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.