

Anti-JIP3 Antibody

Rabbit polyclonal antibody to JIP3

Catalog # AP59816

Product Information

Application	WB, IF/IC, IHC
Primary Accession	Q9UPT6
Other Accession	Q9ESN9
Reactivity	Human, Mouse, Rat, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	147457

Additional Information

Gene ID	23162
Other Names	JIP3; KIAA1066; C-Jun-amino-terminal kinase-interacting protein 3; JIP-3; JNK-interacting protein 3; JNK MAP kinase scaffold protein 3; Mitogen-activated protein kinase 8-interacting protein 3
Target/Specificity	Recognizes endogenous levels of JIP3 protein.
Dilution	WB~~WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500) IF/IC~~N/A IHC~~WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	MAPK8IP3
Synonyms	JIP3, KIAA1066
Function	The JNK-interacting protein (JIP) group of scaffold proteins selectively mediates JNK signaling by aggregating specific components of the MAPK cascade to form a functional JNK signaling module (PubMed: 12189133). May function as a regulator of vesicle transport, through interactions with the JNK-signaling components and motor proteins (By similarity). Promotes neuronal axon elongation in a kinesin- and JNK-dependent manner. Activates cofilin at axon tips via local activation of JNK, thereby regulating filopodial dynamics and enhancing axon elongation. Its binding to kinesin heavy chains (KHC), promotes kinesin-1 motility along microtubules and is essential for

axon elongation and regeneration. Regulates cortical neuronal migration by mediating NTRK2/TRKB anterograde axonal transport during brain development (By similarity). Acts as an adapter that bridges the interaction between NTRK2/TRKB and KLC1 and drives NTRK2/TRKB axonal but not dendritic anterograde transport, which is essential for subsequent BDNF-triggered signaling and filopodia formation (PubMed:[21775604](https://pubmed.ncbi.nlm.nih.gov/21775604/)).


Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:Q9ESN9}. Golgi apparatus {ECO:0000250|UniProtKB:Q9ESN9}. Cytoplasmic vesicle {ECO:0000250|UniProtKB:Q9ESN9}. Cell projection, growth cone {ECO:0000250|UniProtKB:Q9ESN9}. Cell projection, axon {ECO:0000250|UniProtKB:Q9ESN9}. Cell projection, dendrite {ECO:0000250|UniProtKB:E9PSK7}. Cytoplasm, perinuclear region {ECO:0000250|UniProtKB:E9PSK7}. Note=Localized in the soma and growth cones of differentiated neurites and the Golgi and vesicles of the early secretory compartment of epithelial cells. KIF5A/B/C-mediated transportation to axon tips is essential for its function in enhancing neuronal axon elongation. {ECO:0000250|UniProtKB:E9PSK7, ECO:0000250|UniProtKB:Q9ESN9}


Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human JIP3. The exact sequence is proprietary.

Images




Western blot analysis of JIP3 expression in BV2 (A), HuT78 (B), Hela (C), HEK293T (D) whole cell lysates.



Immunohistochemical analysis of JIP3 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Immunofluorescent analysis of JIP3 staining in PC12 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and



incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.