10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-DDX54 Antibody

Catalog # AP59870

Product Information

Application	WB
Primary Accession	Q8TDD1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	98595

Additional Information


Gene ID	79039
Other Names	ATP-dependent RNA helicase DDX54; ATP-dependent RNA helicase DP97; DEAD box RNA helicase 97 kDa; DEAD box protein 54
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human DDX54. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000), IP (1/10 - 1/100)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	DDX54
Function	Nucleolar RNA helicase that is involved in diverse cellular processes including ribosome biogenesis, DNA damage response, RNA splicing, transcriptional regulation, innate immunity, and nervous system development. Plays an essential role in the formation of the ribosome active site by remodeling rRNA structure and initiating peptidyl transferase center formation (PubMed:38632236). Represses the transcriptional activity of several nuclear receptors. Regulates transcriptome dynamics during DNA damage response. Mechanistically, acts via increased interaction with a well-defined class of pre-mRNAs that contain introns with weak acceptor splice sites, as well as by protein-protein contacts within components of U2 snRNP and spliceosomal B complex. These activities lead to reduced intron retention and enhanced processing rates of its target transcripts (PubMed:28596291). Acts as an inhibitor of type I interferon antiviral response by facilitating ALKBH5-mediated demethylation of specific transcripts (PubMed:40793791). Contributes to central nervous system

	myelination, likely through regulation of oligodendrocyte differentiation and myelin sheath formation (By similarity).
Cellular Location	Nucleus, nucleolus. Nucleus, nucleoplasm Note=Relocates from the nucleolus to the nucleoplasm upon viral infection.
Tissue Location	Expressed in the fallopian tube, cervix and uterus. Also expressed in the brain.

Images



Western blot analysis of DDX54 expression in H446 (A) whole cell lysates. (Predicted band size: 98 kD; Observed band size: 98 kD)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.