

Anti-NAA15 Antibody

Catalog # AP59875

Product Information

Application	WB, IHC, IF/IC
Primary Accession	Q9BXJ9
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	101272

Additional Information

Gene ID	80155
Other Names	GA19; NARG1; NATH; TBDN100; N-alpha-acetyltransferase 15, NatA auxiliary subunit; Gastric cancer antigen Ga19; N-terminal acetyltransferase; NMDA receptor-regulated protein 1; Protein tubedown-1; Tbdn100
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human NAA15. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500) IHC~~WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500) IF/IC~~N/A
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt


Protein Information

Name	NAA15
Synonyms	GA19, NARG1, NATH, TBDN100
Function	Auxillary subunit of N-terminal acetyltransferase complexes which display alpha (N-terminal) acetyltransferase (NAT) activity (PubMed: 15496142 , PubMed: 20154145 , PubMed: 29754825 , PubMed: 32042062). The NAT activity may be important for vascular, hematopoietic and neuronal growth and development (PubMed: 15496142). Required to control retinal neovascularization in adult ocular endothelial cells (PubMed: 11687548). In complex with XRCC6 and XRCC5 (Ku80), up-regulates transcription from the osteocalcin promoter (PubMed: 12145306).
Cellular Location	Cytoplasm. Nucleus. Note=Mainly cytoplasmic, nuclear in some cases. Present in the free cytosolic and cytoskeleton- bound polysomes, but not in the


Tissue Location

Expressed at high levels in testis and in ocular endothelial cells. Also found in brain (corpus callosum), heart, colon, bone marrow and at lower levels in most adult tissues, including thyroid, liver, pancreas, mammary and salivary glands, lung, ovary, urogenital system and upper gastrointestinal tract. Overexpressed in gastric cancer, in papillary thyroid carcinomas and in a Burkitt lymphoma cell line (Daudi). Specifically suppressed in abnormal proliferating blood vessels in eyes of patients with proliferative diabetic retinopathy.


Images



Western blot analysis of NAA15 expression in A549 (A), DLD (B), H446 (C), mouse muscle (D), mouse testis (E), rat muscle (F) whole cell lysates. (Predicted band size: 101 kD; Observed band size: 101 kD)




Immunohistochemical analysis of NAA15 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of NAA15 staining in THP1 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

Immunohistochemical analysis of NAA15 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated



compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.