

Anti-CTIP (pS327) Antibody

Rabbit polyclonal antibody to CTIP (pS327)

Catalog # AP59933

Product Information

Application	WB
Primary Accession	Q99708
Other Accession	Q80YR6
Reactivity	Human, Mouse, Monkey
Host	Rabbit
Clonality	Polyclonal
Calculated MW	101942

Additional Information

Gene ID	5932
Other Names	CTIP; DNA endonuclease RBBP8; CtBP-interacting protein; CtIP; Retinoblastoma-binding protein 8; RBBP-8; Retinoblastoma-interacting protein and myosin-like; RIM; Sporulation in the absence of SPO11 protein 2 homolog; SAE2
Target/Specificity	Recognizes endogenous levels of CTIP (pS327) protein.
Dilution	WB~~WB (1/500 - 1/1000)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	RBBP8
Synonyms	CTIP
Function	Endonuclease that cooperates with the MRE11-RAD50-NBN (MRN) complex in DNA-end resection, the first step of double-strand break (DSB) repair through the homologous recombination (HR) pathway (PubMed: 17965729 , PubMed: 19202191 , PubMed: 19759395 , PubMed: 20064462 , PubMed: 23273981 , PubMed: 26721387 , PubMed: 27814491 , PubMed: 27889449 , PubMed: 30787182). HR is restricted to S and G2 phases of the cell cycle and preferentially repairs DSBs resulting from replication fork collapse (PubMed: 17965729 , PubMed: 19202191 , PubMed: 23273981 , PubMed: 27814491 , PubMed: 27889449 , PubMed: 30787182). Key determinant of DSB repair pathway choice, as it commits cells to HR by preventing classical

non-homologous end-joining (NHEJ) (PubMed:[19202191](#)). Specifically promotes the endonuclease activity of the MRN complex to clear DNA ends containing protein adducts: recruited to DSBs by NBN following phosphorylation by CDK1, and promotes the endonuclease activity of MRE11 to clear protein-DNA adducts and generate clean double-strand break ends (PubMed:[27814491](#), PubMed:[27889449](#), PubMed:[30787182](#), PubMed:[33836577](#)). Functions downstream of the MRN complex and ATM, promotes ATR activation and its recruitment to DSBs in the S/G2 phase facilitating the generation of ssDNA (PubMed:[16581787](#), PubMed:[17965729](#), PubMed:[19759395](#), PubMed:[20064462](#)). Component of the BRCA1-RBBP8 complex that regulates CHEK1 activation and controls cell cycle G2/M checkpoints on DNA damage (PubMed:[15485915](#), PubMed:[16818604](#)). During immunoglobulin heavy chain class-switch recombination, promotes microhomology-mediated alternative end joining (A-NHEJ) and plays an essential role in chromosomal translocations (By similarity). Binds preferentially to DNA Y-junctions and to DNA substrates with blocked ends and promotes intermolecular DNA bridging (PubMed:[30601117](#)).

Cellular Location

Nucleus. Chromosome Note=Associates with sites of DNA damage in S/G2 phase (PubMed:[10764811](#), PubMed:[25349192](#)). Recruited to DSBs by the MRE11- RAD50-NBN (MRN) complex following phosphorylation by CDK1, which promotes interaction with NBN (PubMed:[27814491](#), PubMed:[27889449](#), PubMed:[33836577](#)). Ubiquitinated RBBP8 binds to chromatin following DNA damage (PubMed:[16818604](#)).


Tissue Location

Expressed in ER-positive breast cancer lines, but tends to be down-regulated ER-negative cells (at protein level)


Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human CTIP (pS327). The exact sequence is proprietary.

Images



Western blot analysis of CTIP (pS327) expression in U87 (A), PC3 (B), MCF7 (C) whole cell lysates.



Direct ELISA antibody dose-response curve using Anti-CTIP (pS327) Antibody. Antigen (phosphopeptide and non-phosphopeptide) concentration is 5 ug/ml. Goat Anti-Rabbit IgG (H&L) - HRP was used as the secondary antibody, and signal was developed by TMB substrate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.