10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-MINA53 Antibody

Rabbit polyclonal antibody to MINA53
Catalog # AP59954

Product Information

Application	WB
Primary Accession	Q8IUF8
Other Accession	Q8CD15
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	52800

Additional Information

Gene ID	84864
Other Names	MDIG; MINA53; NO52; Bifunctional lysine-specific demethylase and histidyl-hydroxylase MINA; 60S ribosomal protein L27a histidine hydroxylase; Histone lysine demethylase MINA; MYC-induced nuclear antigen; Mineral dust-induced gene protein; Nucleolar protein 52; Ribosomal oxygenase MINA; ROX
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human MINA53. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	RIOX2 (HGNC:19441)
Function	Oxygenase that can act as both a histone lysine demethylase and a ribosomal histidine hydroxylase. Is involved in the demethylation of trimethylated 'Lys-9' on histone H3 (H3K9me3), leading to an increase in ribosomal RNA expression. Also catalyzes the hydroxylation of 60S ribosomal protein L27a on 'His-39'. May play an important role in cell growth and survival. May be involved in ribosome biogenesis, most likely during the assembly process of pre-ribosomal particles.
Cellular Location	Nucleus. Nucleus, nucleolus

Tissue Location	Expressed in liver, skeletal muscle, heart, pancreas, and placenta. Not detected in brain, lung or kidney Expressed in several lung cancer tissues, but is barely detected in the adjacent non-cancerous tissues. Also highly expressed in several esophageal squamous cell carcinoma (ESCC), and colon cancer tissues, and in various cancer cell lines.

Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human MINA53. The exact sequence is proprietary.

Images



Western blot analysis of MINA53 expression in rat liver (A) whole cell lysates.