

# Anti-Androgen Receptor (pS650) Antibody

Rabbit polyclonal antibody to Androgen Receptor (pS650)

Catalog # AP59958

## Product Information

Application	WB, IHC
Primary Accession	<a href="#">P10275</a>
Other Accession	<a href="#">P19091</a>
Reactivity	Human, Mouse, Rat, Monkey
Host	Rabbit
Clonality	Polyclonal
Calculated MW	99188

## Additional Information

Gene ID	367
Other Names	DHTR; NR3C4; Androgen receptor; Dihydrotestosterone receptor; Nuclear receptor subfamily 3 group C member 4
Target/Specificity	Recognizes endogenous levels of Androgen Receptor (pS650) protein.
Dilution	WB~~WB (1/500 - 1/1000), IHC (1/100 - 1/500) IHC~~WB (1/500 - 1/1000), IHC (1/100 - 1/500)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

## Protein Information

Name	AR
Synonyms	DHTR, NR3C4
Function	<p>Steroid hormone receptors are ligand-activated transcription factors that regulate eukaryotic gene expression and affect cellular proliferation and differentiation in target tissues (PubMed:<a href="#">19022849</a>). Transcription factor activity is modulated by bound coactivator and corepressor proteins like ZBTB7A that recruits NCOR1 and NCOR2 to the androgen response elements/ARE on target genes, negatively regulating androgen receptor signaling and androgen-induced cell proliferation (PubMed:<a href="#">20812024</a>). Transcription activation is also down-regulated by NR0B2. Activated, but not phosphorylated, by HIPK3 and ZIPK/DAPK3.</p> <p>Nucleus. Cytoplasm Note=Detected at the promoter of target genes</p>

## Cellular Location

(PubMed:25091737) Predominantly cytoplasmic in unligated form but translocates to the nucleus upon ligand-binding. Can also translocate to the nucleus in unligated form in the presence of RACK1.


## Tissue Location

[Isoform 2]: Mainly expressed in heart and skeletal muscle.


## Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Androgen Receptor (pS650). The exact sequence is proprietary.


## Images



Western blot analysis of Androgen Receptor (pS650) expression in H1688 (A), mouse liver (B), mouse kidney (C), rat liver (D), rat kidney (E) whole cell lysates.



Immunohistochemical analysis of Androgen Receptor (pS650) staining in human prostate cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Direct ELISA antibody dose-response curve using Anti-Androgen Receptor (pS650) Antibody. Antigen (phosphopeptide and non-phosphopeptide) concentration is 5 ug/ml. Goat Anti-Rabbit IgG (H&L) - HRP was used as the secondary antibody, and signal was developed by TMB substrate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.