

Anti-HEXB Antibody

Rabbit polyclonal antibody to HEXB
Catalog # AP60011

Product Information

Application	WB, IF/IC, IHC
Primary Accession	P07686
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	63137

Additional Information

Gene ID	3074
Other Names	Beta-hexosaminidase subunit beta; Beta-N-acetylhexosaminidase subunit beta; Hexosaminidase subunit B; Cervical cancer proto-oncogene 7 protein; HCC-7; N-acetyl-beta-glucosaminidase subunit beta
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human HEXB. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500) IF/IC~~N/A IHC~~WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C. Stable for 12 months from date of receipt

Protein Information

Name	HEXB (HGNC:4879)
Function	Hydrolyzes the non-reducing end N-acetyl-D-hexosamine and/or sulfated N-acetyl-D-hexosamine of glycoconjugates, such as the oligosaccharide moieties from proteins and neutral glycolipids, or from certain mucopolysaccharides (PubMed: 11707436 , PubMed: 8123671 , PubMed: 8672428 , PubMed: 9694901). The isozyme B does not hydrolyze each of these substrates, however hydrolyzes efficiently neutral oligosaccharide (PubMed: 11707436). Only the isozyme A is responsible for the degradation of GM2 gangliosides in the presence of GM2A (PubMed: 8123671 , PubMed: 8672428 , PubMed: 9694901). During fertilization is responsible, at least in part, for the zona block to polyspermy. Present in the cortical granules of non-activated oocytes, is exocytosed during the cortical reaction in

response to oocyte activation and inactivates the sperm galactosyltransferase-binding site, accounting for the block in sperm binding to the zona pellucida (By similarity).


Cellular Location

Lysosome. Cytoplasmic vesicle, secretory vesicle, Cortical granule
{ECO:0000250 | UniProtKB:P20060}


Background

KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human HEXB. The exact sequence is proprietary.


Images



Western blot analysis of HEXB expression in HEK293T (A), A375 (B), LO2 (C) whole cell lysates.



Immunohistochemical analysis of HEXB staining in human prostate cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of HEXB staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.