10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-Kv9.2 Antibody

Catalog # AP60019

Product Information

Application	WB, IF/IC
Primary Accession	Q9ULS6
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	54237

Additional Information


Gene ID	3788
Other Names	KIAA1144; Potassium voltage-gated channel subfamily S member 2; Delayed-rectifier K(+) channel alpha subunit 2; Voltage-gated potassium channel subunit Kv9.2
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Kv9.2. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) IF/IC~~N/A
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information


Name	KCNS2 (HGNC:6301)
Synonyms	KIAA1144
Function	Potassium channel regulatory subunit that modulate the delayed rectifier voltage-gated potassium channel activity of KCNB1 and KCNB2 by altering their kinetics, expression levels, and shifting the half-inactivation potential to more polarized values. While it does not form functional channels on its own, it can form functional heterotetrameric channels with KCNB1 and KCNB2. Each regulatory subunit has unique regulatory properties that can lead to extensive inhibition, significant changes in kinetics, and/or substantial shifts in the voltage dependencies of the inactivation process.
Cellular Location	Cell membrane {ECO:0000250|UniProtKB:O35174}; Multi-pass membrane protein {ECO:0000250|UniProtKB:O35174}. Note=May not reach the plasma membrane but remain in an intracellular compartment in the absence of

KCNB1 or KCNB2 {ECO:0000250|UniProtKB:O35174}

Images



Western blot analysis of Kv9.2 expression in A2780 (A) whole cell lysates. (Predicted band size: 54 kD; Observed band size: 54 kD)



Immunofluorescent analysis of Kv9.2 staining in SGC7901 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with a AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.