10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-LKB1 Antibody

Catalog # AP60058

Product Information

Application	WB
Primary Accession	Q15831
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	48636

Additional Information

Gene ID	6794
Other Names	LKB1; PJS; Serine/threonine-protein kinase STK11; Liver kinase B1; LKB1; hLKB1; Renal carcinoma antigen NY-REN-19
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human LKB1. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	STK11 (HGNC:11389)
Synonyms	LKB1, PJS
Function	Tumor suppressor serine/threonine-protein kinase that controls the activity of AMP-activated protein kinase (AMPK) family members, thereby playing a role in various processes such as cell metabolism, cell polarity, apoptosis and DNA damage response (PubMed:12805220, PubMed:14517248, PubMed:14976552, PubMed:15016379, PubMed:15733851). Acts by phosphorylating the T-loop of AMPK family proteins, thus promoting their activity: phosphorylates PRKAA1, PRKAA2, BRSK1, BRSK2, MARK1, MARK2, MARK3, MARK4, NUAK1, NUAK2, SIK1, SIK2, SIK3 and SNRK but not MELK (PubMed:12805220, PubMed:14517248, PubMed:14976552, PubMed:15016379, PubMed:15733851). Also phosphorylates non-AMPK family proteins such as STRADA, PTEN and possibly p53/TP53 (PubMed:11430832, PubMed:15987703). Acts as a key upstream regulator of AMPK by mediating phosphorylation and activation of AMPK catalytic subunits

PRKAA1 and PRKAA2 and thereby regulates processes including: inhibition of signaling pathways that promote cell growth and proliferation when energy levels are low, glucose homeostasis in liver, activation of autophagy when cells undergo nutrient deprivation, and B- cell differentiation in the germinal center in response to DNA damage (PubMed:14976552). Also acts as a regulator of cellular polarity by remodeling the actin cytoskeleton (PubMed:15016379). Required for cortical neuron polarization by mediating phosphorylation and activation of BRSK1 and BRSK2, leading to axon initiation and specification (PubMed:14976552). Involved in DNA damage response: interacts with p53/TP53 and recruited to the CDKN1A/WAF1 promoter to participate in transcription activation (PubMed:11430832, PubMed:17108107, PubMed:21317932). Able to phosphorylate p53/TP53; the relevance of such result in vivo is however unclear and phosphorylation may be indirect and mediated by downstream STK11/LKB1 kinase NUAK1 (PubMed:11430832, PubMed:17108107, PubMed:21317932). Also acts as a mediator of p53/TP53-dependent apoptosis via interaction with p53/TP53: translocates to the mitochondrion during apoptosis and regulates p53/TP53-dependent apoptosis pathways (PubMed:11430832, PubMed:17108107, PubMed:21317932). Regulates UV radiation-induced DNA damage response mediated by CDKN1A (PubMed:25329316). In association with NUAK1, phosphorylates CDKN1A in response to UV radiation and contributes to its degradation which is necessary for optimal DNA repair (PubMed:25329316).


Cellular Location

Nucleus. Cytoplasm. Membrane {ECO:0000250|UniProtKB:Q9WTK7} Mitochondrion. Late endosome {ECO:0000250|UniProtKB:Q9WTK7}. Note=A small fraction localizes at membranes (By similarity). Relocates to the cytoplasm when bound to STRAD (STRADA or STRADB) and CAB39/MO25 (CAB39/MO25alpha or CAB39L/MO25beta) (PubMed:18854309). Translocates to the mitochondrion during apoptosis. PTEN promotes cytoplasmic localization (PubMed:11430832). Translocates to late endosomes in response to calorie restriction (By similarity). {ECO:0000250|UniProtKB:Q9WTK7, ECO:0000269|PubMed:11430832, ECO:0000269|PubMed:18854309}

Tissue Location

Ubiquitously expressed. Strongest expression in testis and fetal liver

Images



Western blot analysis of LKB1 expression in rat testis (A) whole cell lysates. (Predicted band size: 48 kD; Observed band size: 54 kD)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.