

Anti-Histone H4 (pS47) Antibody

Rabbit polyclonal antibody to Histone H4 (pS47)

Catalog # AP60064

Product Information

| | |
|-------------------|------------------------|
| Application | WB, IF/IC |
| Primary Accession | P62805 |
| Reactivity | Human, Rat |
| Host | Rabbit |
| Clonality | Polyclonal |
| Calculated MW | 11367 |

Additional Information

| | |
|--------------------|---|
| Gene ID | 121504;554313;8294;8359;8360;8361;8362;8363;8364;8365;8366;8367;8368;8370 |
| Other Names | H4/A; H4FA; H4/I; H4FI; H4/G; H4FG; H4/B; H4FB; H4/J; H4FJ; H4/C; H4FC; H4/H; H4FH; H4/M; H4FM; H4/E; H4FE; H4/D; H4FD; H4/K; H4FK; H4/N; H4F2; H4FN; HIST2H4; H4/O; H4FO; Histone H4 |
| Target/Specificity | KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Histone H4 (pS47). The exact sequence is proprietary. |
| Dilution | WB~~WB (1/500 - 1/1000), IF/IC (1/100 - 1/500) IF/IC~~N/A |
| Format | Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide. |
| Storage | Store at -20 °C.Stable for 12 months from date of receipt |


Protein Information

| | |
|-------------------|--|
| Name | H4C1 |
| Synonyms | H4/A, H4FA, HIST1H4A |
| Function | Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling. |
| Cellular Location | Nucleus {ECO:0000250 UniProtKB:P62806}. Chromosome. Note=Localized to the nucleus when acetylated in step 11 spermatids. {ECO:0000250 UniProtKB:P62806} |


Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Histone H4 (pS47). The exact sequence is proprietary.

Images



Western blot analysis of Histone H4 (pS47) expression in HEK293T (A), U2OS (B) whole cell lysates.



Immunofluorescent analysis of Histone H4 (pS47) staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.