

SARS virus PUP6 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP6006b

Product Information

Application	E
Primary Accession	P59636
Reactivity	SARS
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB3807-3808
Calculated MW	10802
Antigen Region	69-98

Additional Information

Other Names	Protein 9b, Accessory protein 9b, ORF-9b, 9b
Target/Specificity	This SARS virus PUP6 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 69~98 amino acids from the C-terminal region of human SARS virus protein PUP6.
Dilution	E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	SARS virus PUP6 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	9b
Function	Plays a role in the inhibition of host innate immune response by targeting the mitochondrial-associated adapter MAVS. Mechanistically, usurps the E3 ligase ITCH to trigger the degradation of MAVS, TRAF3, and TRAF6. In addition, causes mitochondrial elongation by triggering ubiquitination and proteasomal degradation of dynamin- like protein 1/DNM1L.
Cellular Location	Virion. Host cytoplasmic vesicle membrane; Peripheral membrane protein.

Host cytoplasm. Host endoplasmic reticulum. Host nucleus. Host mitochondrion. Note=Binds non-covalently to intracellular lipid bilayers

Background

An outbreak of atypical pneumonia, referred to as severe acute respiratory syndrome (SARS) and first identified in Guangdong Province, China, has spread to several countries. The severity of this disease is such that the mortality rate appears to be ~3 to 6%. A number of laboratories worldwide have undertaken the identification of the causative agent. The National Microbiology Laboratory in Canada obtained the Tor2 isolate from a patient in Toronto, and succeeded in growing a coronavirus-like agent in African Green Monkey Kidney (Vero E6) cells. This coronavirus has been named publicly by the World Health Organization and member laboratories as ?SARS virus? The SARS membrane proteins, including the major proteins S (Spike) and M (Membrane), are inserted into the endoplasmic reticulum Golgi intermediate compartment (ERGIC) while full length replicated RNA (+ strands) assemble with the N (nucleocapsid) protein. The virus then migrates through the Golgi complex and eventually exits the cell, likely by exocytosis. The site of viral attachment to the host cell resides within the S protein. Oligomeric spike (S) glycoproteins extend from SARS membranes. These integral membrane proteins assemble within the endoplasmic reticulum of infected cells and are subsequently endoproteolyzed in the Golgi, generating noncovalently associated S1 and S2 fragments. Once on the surface of infected cells and virions, peripheral S1 fragments bind carcinoembryonic antigen-related cell adhesion molecule (CEACAM) receptors, and this triggers membrane fusion reactions mediated by integral membrane S2 fragments.

References

He, R., et al., Biochem. Biophys. Res. Commun. 316(2):476-483 (2004).
Snijder, E.J., et al., J. Mol. Biol. 331(5):991-1004 (2003).
Marra, M.A., et al., Science 300(5624):1399-1404 (2003).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.