

# Anti-TFCP2L1 Antibody

Rabbit polyclonal antibody to TFCP2L1

Catalog # AP60206

## Product Information

---

Application	WB
Primary Accession	<a href="#">Q9NZI6</a>
Other Accession	<a href="#">Q3UNW5</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	54627

## Additional Information

---

Gene ID	29842
Other Names	CRTR1; LBP9; Transcription factor CP2-like protein 1; CP2-related transcriptional repressor 1; CRTR-1; Transcription factor LBP-9
Target/Specificity	Recognizes endogenous levels of TFCP2L1 protein.
Dilution	WB~~WB (1/500 - 1/1000)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

## Protein Information

---

Name	TFCP2L1
Synonyms	CRTR1, LBP9
Function	Transcription factor that facilitates establishment and maintenance of pluripotency in embryonic stem cells (ESCs) (PubMed: <a href="#">25215486</a> , PubMed: <a href="#">26906118</a> ). With KLF2, acts as the major effector of self-renewal that mediates induction of pluripotency downstream of LIF/STAT3 and Wnt/beta-catenin signaling (By similarity). Required for normal duct development in the salivary gland and kidney (By similarity). Coordinates the development of the kidney collecting ducts intercalated (IC) and principal (PC) cells, which regulate acid- base and salt-water homeostasis, respectively (By similarity). Regulates the expression of IC genes including subunits B1 and D2 of the V-ATPase complex, OXGR1, CA12, SLC4A1, AQP6 and IC-specific transcription factor FOXI1 (By similarity). Also regulates the expression of JAG1 and subsequent notch signaling in the collecting duct (By similarity).

JAG1 initiates notch signaling in PCs but inhibits notch signaling in ICs (By similarity). Acts as a transcriptional suppressor that may suppress UBPI-mediated transcriptional activation (By similarity). Modulates the placental expression of CYP11A1 (PubMed:[10644752](#)).

**Cellular Location**

Nucleus.

**Tissue Location**

Highly expressed in placental JEG-3 cells and very low levels of expression in non-steroidogenic cells. No expression was seen in adrenal NCI-H295A cells or in adrenal tissue


## Background

---

KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human TFCP2L1. The exact sequence is proprietary.

## Images

---



Western blot analysis of TFCP2L1 expression in HEK293T (A) whole cell lysates.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.