

# Anti-CD274 Antibody

Rabbit polyclonal antibody to CD274 Catalog # AP60240

### **Product Information**

Application WB
Primary Accession Q9NZQ7
Other Accession Q9EP73

**Reactivity** Human, Mouse, Rat

Host Rabbit
Clonality Polyclonal
Calculated MW 33275

## **Additional Information**

**Gene ID** 29126

Other Names B7H1; PDCD1L1; PDCD1LG1; PDL1; Programmed cell death 1 ligand 1; PD-L1;

PDCD1 ligand 1; Programmed death ligand 1; B7 homolog 1; B7-H1; CD274

**Target/Specificity** KLH-conjugated synthetic peptide encompassing a sequence within the center

region of human CD274. The exact sequence is proprietary.

**Dilution** WB~~WB (1/500 - 1/1000)

**Format** Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30%

glycerol, and 0.09% (W/V) sodium azide.

**Storage** Store at -20 °C.Stable for 12 months from date of receipt

## **Protein Information**

Name CD274 ( <u>HGNC:17635</u>)

**Function** Plays a critical role in induction and maintenance of immune tolerance to

self (PubMed:<u>11015443</u>, PubMed:<u>28813410</u>, PubMed:<u>28813417</u>,

PubMed:31399419). As a ligand for the inhibitory receptor PDCD1/PD-1, modulates the activation threshold of T-cells and limits T-cell effector response (PubMed:11015443, PubMed:28813410, PubMed:28813417, PubMed:36727298). Through a yet unknown activating receptor, may costimulate T-cell subsets that predominantly produce interleukin-10 (IL10) (PubMed:10581077). Can also act as a transcription coactivator: in response

to hypoxia, translocates into the nucleus via its interaction with

phosphorylated STAT3 and promotes transcription of GSDMC, leading to

pyroptosis (PubMed:32929201).

Cell membrane; Single-pass type I membrane protein. Early endosome

#### **Cellular Location**

membrane; Single-pass type I membrane protein. Recycling endosome membrane; Single-pass type I membrane protein. Nucleus. Note=Associates with CMTM6 at recycling endosomes, where it is protected from being targeted for lysosomal degradation (PubMed:28813417). Translocates to the nucleus in response to hypoxia via its interaction with phosphorylated STAT3 (PubMed:32929201). [Isoform 2]: Endomembrane system; Single-pass type I membrane protein


#### **Tissue Location**

Highly expressed in the heart, skeletal muscle, placenta and lung. Weakly expressed in the thymus, spleen, kidney and liver. Expressed on activated T-and B-cells, dendritic cells, keratinocytes and monocytes.

# **Background**

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human CD274. The exact sequence is proprietary.

# **Images**



Western blot analysis of CD274 expression in mouse lung (A) whole cell lysates.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.