10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-FRS2 (Phospho-Y436) Antibody

Catalog # AP60287

Product Information

Application	WB, IHC
Primary Accession	Q8WU20
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	57029


Additional Information

Gene ID	10818
Other Names	Fibroblast growth factor receptor substrate 2; FGFR substrate 2; FGFR-signaling adaptor SNT; Suc1-associated neurotrophic factor target 1; SNT-1
Target/Specificity	KLH-conjugated synthetic phosphopeptide corresponding to residues surrounding Y436 of human FRS2 protein. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000), IHC (1/100 - 1/200) IHC~~WB (1/500 - 1/1000), IHC (1/100 - 1/200)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt


Protein Information

Name	FRS2
Function	Adapter protein that links activated FGR and NGF receptors to downstream signaling pathways. Plays an important role in the activation of MAP kinases and in the phosphorylation of PIK3R1, the regulatory subunit of phosphatidylinositol 3-kinase, in response to ligand-mediated activation of FGFR1. Modulates signaling via SHC1 by competing for a common binding site on NTRK1.
Cellular Location	Endomembrane system. Note=Cytoplasmic, membrane- bound
Tissue Location	Highly expressed in heart, brain, spleen, lung, liver, skeletal muscle, kidney and testis


Images



Western blot analysis of FRS2 (Phospho-Y436) expression in U87MG (A), Hela (B), PC12 (C) whole cell lysates. (Predicted band size: 57 kD; Observed band size: 57; 75 kD)



Immunohistochemical analysis of FRS2 (Phospho-Y436) staining in human liver formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunohistochemical analysis of FRS2 (pY436) staining in human liver formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.