

## Anti-GBA3 Antibody

Rabbit polyclonal antibody to GBA3  
Catalog # AP60292

### Product Information

---

<b>Application</b>	WB
<b>Primary Accession</b>	<a href="#">Q9H227</a>
<b>Reactivity</b>	Human, Mouse, Rat
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	53696

### Additional Information

---

<b>Gene ID</b>	57733
<b>Other Names</b>	CBG; CBGL1; Cytosolic beta-glucosidase; Cytosolic beta-glucosidase-like protein 1
<b>Target/Specificity</b>	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human GBA3. The exact sequence is proprietary.
<b>Dilution</b>	WB~~WB (1/500 - 1/1000)
<b>Format</b>	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
<b>Storage</b>	Store at -20 °C. Stable for 12 months from date of receipt

### Protein Information

---

<b>Name</b>	GBA3 ( <a href="#">HGNC:19069</a> )
<b>Synonyms</b>	CBG, CBGL1
<b>Function</b>	Neutral cytosolic beta-glycosidase with a broad substrate specificity that could play a role in the catabolism of glycosylceramides (PubMed: <a href="#">11389701</a> , PubMed: <a href="#">11784319</a> , PubMed: <a href="#">17595169</a> , PubMed: <a href="#">20728381</a> , PubMed: <a href="#">26724485</a> , PubMed: <a href="#">33361282</a> ). Has a significant glucosylceramidase activity in vitro (PubMed: <a href="#">17595169</a> , PubMed: <a href="#">26724485</a> ). However, that activity is relatively low and its significance in vivo is not clear (PubMed: <a href="#">17595169</a> , PubMed: <a href="#">20728381</a> , PubMed: <a href="#">26724485</a> ). Hydrolyzes galactosylceramides/GalCers, glucosylsphingosines/GlcSphs and galactosylsphingosines/GalSphs (PubMed: <a href="#">17595169</a> ). However, the in vivo relevance of these activities is unclear (PubMed: <a href="#">17595169</a> ). It can also hydrolyze a broad variety of dietary glycosides including phytoestrogens, flavonols, flavones, flavanones and cyanogens in vitro and could therefore

play a role in the metabolism of xenobiotics (PubMed:[11784319](#)). Possesses transxylosylase activity in vitro using xylosylated ceramides/XylCers (such as beta-D-xylosyl-(11')-N-acylsphing-4-enine) as xylosyl donors and cholesterol as acceptor (PubMed:[33361282](#)). Could also play a role in the catabolism of cytosolic sialyl free N-glycans (PubMed:[26193330](#)).

#### Cellular Location

Cytoplasm, cytosol

#### Tissue Location

Present in small intestine (at protein level). Expressed in liver, small intestine, colon, spleen and kidney. Down- regulated in renal cell carcinomas and hepatocellular carcinomas


## Background

---

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human GBA3. The exact sequence is proprietary.

## Images

---



Western blot analysis of GBA3 expression in LO2 (A), HepG2 (B), CT26 (C), rat liver (D) whole cell lysates.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.