10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-LIF Antibody

Catalog # AP60332

Product Information

Application	WB, IHC, IF/IC
Primary Accession	P15018
Reactivity	Human, Mouse, Rat, Pig, Bovine, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	22008

Additional Information

Gene ID	3976
Other Names	HILDA; Leukemia inhibitory factor; LIF; Differentiation-stimulating factor; D factor; Melanoma-derived LPL inhibitor; MLPLI; Emfilermin
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human LIF. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000), IHC (1/50 - 1/200) IHC~~WB (1/500 - 1/1000), IHC (1/50 - 1/200) IF/IC~~N/A
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt


Protein Information

Name	LIF (HGNC:6596)
Synonyms	HILDA
Function	Functions as a cytokine that binds to the LIF receptor complex (PubMed:11294841, PubMed:14527405, PubMed:36930708). The LIF receptor complex indeed consists of two signaling receptor subunits IL6ST/gp130 and LIFR that transduce the signal into the cell (PubMed:14527405, PubMed:36930708). Functionally, regulates cell self- renewal, differentiation and survival of embryonic stem cells (ESCs) and is crucial for embryonic implantation (By similarity). Mechanistically, ligand binding to LIFR, induces heterodimerization with IL6ST/gp130 activating JAK tyrosine kinases (JAK1 or JAK2 and to a lesser extent TYK2) bound to their intracellular domains (PubMed:11294841). These kinases subsequently phosphorylate IL6ST/gp130 and LIFR (PubMed:11294841). The tyrosine phosphorylated signaling receptors serve in turn as docking sites for recruitment and activation of


signal transducers and activators of transcription (STAT3 and to lesser extent STAT1) (PubMed:11294841). In parallel to the activation of the STAT3-pathway, the binding of LIF to the LIF receptor complex leads to the activation of the mitogen-activated protein kinase (MAPK) and the phosphatidylinositol 3-kinase (PI(3)K) pathways (PubMed:11294841). Tyrosine phosphorylation of the p85 subunit of PI(3)K by JAK1 leads to activation of the serine/ threonine kinase AKT (protein kinase B) (PubMed:11294841). PI(3)K-dependent signaling is required for efficient self-renewal of murine ES cells, and regulation of ERK activity is functionally important in this response (By similarity). Moreover, induces tyrosine phosphorylation of PTPN11, that associates with IL6ST/gp130-LIFR, which in turn recruits the p85 regulatory subunit of PI(3)K protein kinase, promoting PI3K-AKT activation, and GRB2-RAS-MAPK signaling (PubMed:11294841).

Cellular Location Secreted.


Images



Western blot analysis of LIF expression in HCT116 (A), LO2 (B), mouse testis (C), rat testis (D) whole cell lysates. (Predicted band size: 22 kD; Observed band size: 24 kD)




Immunohistochemical analysis of LIF staining in human colon cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of LIF staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with a AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Immunohistochemical analysis of LIF staining in human



colon cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.