

# Anti-p47 phox (pS304) Antibody

Rabbit polyclonal antibody to p47 phox (pS304)

Catalog # AP60344

## Product Information

|                   |                                      |
|-------------------|--------------------------------------|
| Application       | WB, IHC                              |
| Primary Accession | <a href="#">P14598</a>               |
| Reactivity        | Human, Mouse, Rat, Bovine, Dog, SARS |
| Host              | Rabbit                               |
| Clonality         | Polyclonal                           |
| Calculated MW     | 44682                                |

## Additional Information

|                    |   |
|--------------------|---|
| Gene ID            | 653361  |
| Other Names        | NOXO2; SH3PXD1A; Neutrophil cytosol factor 1; NCF-1; 47 kDa autosomal chronic granulomatous disease protein; 47 kDa neutrophil oxidase factor; NCF-47K; Neutrophil NADPH oxidase factor 1; Nox organizer 2; Nox-organizing protein 2; SH3 and PX domain-containing protein 1A; p47-phox |
| Target/Specificity | KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human p47 phox (pS304). The exact sequence is proprietary.   |
| Dilution           | WB~~WB (1/500 - 1/1000), IHC (1/100 - 1/200) IHC~~WB (1/500 - 1/1000), IHC (1/100 - 1/200)  |
| Format             | Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.   |
| Storage            | Store at -20 °C.Stable for 12 months from date of receipt   |

## Protein Information

|          |   |
|----------|---|
| Name     | NCF1 ( <a href="#">HGNC:7660</a> )  |
| Synonyms | NOXO2, SH3PXD1A   |
| Function | Subunit of the phagocyte NADPH oxidase complex that mediates the transfer of electrons from cytosolic NADPH to O <sub>2</sub> to produce the superoxide anion (O <sub>2</sub> <sup>-</sup> ) (PubMed: <a href="#">2547247</a> , PubMed: <a href="#">2550933</a> , PubMed: <a href="#">38355798</a> ). In the activated complex, electrons are first transferred from NADPH to flavin adenine dinucleotide (FAD) and subsequently transferred via two heme molecules to molecular oxygen, producing superoxide through an outer-sphere reaction (PubMed: <a href="#">38355798</a> ). Activation of the NADPH oxidase complex is initiated by the assembly of cytosolic subunits of the NADPH |


oxidase complex with the core NADPH oxidase complex to form a complex at the plasma membrane or phagosomal membrane (PubMed:[38355798](#)). This activation process is initiated by phosphorylation dependent binding of the cytosolic NCF1/p47-phox subunit to the C-terminus of CYBA/p22-phox (PubMed:[12732142](#), PubMed:[19801500](#)).

|                          |   |
|--------------------------|---|
| <b>Cellular Location</b> | Cytoplasm, cytosol. Membrane; Peripheral membrane protein; Cytoplasmic side |
| <b>Tissue Location</b>   | Detected in peripheral blood monocytes and neutrophils (at protein level).  |


## Background

KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human p47 phox (pS304). The exact sequence is proprietary.

## Images



Western blot analysis of p47 phox (pS304) expression in HEK293T (A), Hela (B), mouse lung (C), mouse liver (D), rat liver (E), rat spleen (F) whole cell lysates.



Immunohistochemical analysis of p47 phox (pS304) staining in human lymph node formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.