

# Anti-NFIL3 Antibody

Rabbit polyclonal antibody to NFIL3  
Catalog # AP60348

## Product Information

---

<b>Application</b>	WB, IF/IC
<b>Primary Accession</b>	<a href="#">Q16649</a>
<b>Other Accession</b>	<a href="#">Q08750</a>
<b>Reactivity</b>	Human, Mouse, Rat, Monkey, Chicken, Bovine
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	51472

## Additional Information

---

<b>Gene ID</b>	4783
<b>Other Names</b>	E4BP4; IL3BP1; Nuclear factor interleukin-3-regulated protein; E4 promoter-binding protein 4; Interleukin-3 promoter transcriptional activator; Interleukin-3-binding protein 1; Transcriptional activator NF-IL3A
<b>Target/Specificity</b>	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human NFIL3. The exact sequence is proprietary.
<b>Dilution</b>	WB~~WB (1/500 - 1/1000), IF/IC (1/100 - 1/500) IF/IC~~N/A
<b>Format</b>	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
<b>Storage</b>	Store at -20 °C. Stable for 12 months from date of receipt

## Protein Information

---

<b>Name</b>	NFIL3
<b>Synonyms</b>	E4BP4, IL3BP1
<b>Function</b>	Acts as a transcriptional regulator that recognizes and binds to the sequence 5'-[GA]TTA[CT]GTAA[CT]-3', a sequence present in many cellular and viral promoters. Represses transcription from promoters with activating transcription factor (ATF) sites. Represses promoter activity in osteoblasts (By similarity). Represses transcriptional activity of PER1 (By similarity). Represses transcriptional activity of PER2 via the B-site on the promoter (By similarity). Activates transcription from the interleukin-3 promoter in T-cells. Competes for the same consensus-binding site with PAR DNA-binding factors (DBP, HLF and TEF) (By similarity). Component of the circadian clock that acts as a negative regulator for the circadian expression of PER2 oscillation in the

cell-autonomous core clock (By similarity). Protects pro-B cells from programmed cell death (By similarity). Represses the transcription of CYP2A5 (By similarity). Positively regulates the expression and activity of CES2 by antagonizing the repressive action of NR1D1 on CES2 (By similarity). Required for the development of natural killer cell precursors (By similarity).

#### Cellular Location

Nucleus {ECO:0000255 | PROSITE-ProRule:PRU00978}.


#### Tissue Location

Expressed in bladder, stomach, thyroid, spinal cord, lymph node, trachea, adrenal gland, bone marrow and muscle


## Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human NFIL3. The exact sequence is proprietary.

## Images



Western blot analysis of NFIL3 expression in HeLa (A), mouse lung (B), rat lung (C) whole cell lysates.



Immunofluorescent analysis of NFIL3 staining in K562 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.