

Anti-Phospholipase D4 Antibody

Rabbit polyclonal antibody to Phospholipase D4 Catalog # AP60360

Product Information

Application	WB, IHC
Primary Accession	<u>Q96BZ4</u>
Other Accession	<u>Q8BG07</u>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	55626

Additional Information

Gene ID	122618
Other Names	C14orf175; Phospholipase D4; PLD 4; Choline phosphatase 4; Phosphatidylcholine-hydrolyzing phospholipase D4
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human Phospholipase D4. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000), IHC (1/100 - 1/200) IHC~~WB (1/500 - 1/1000), IHC (1/100 - 1/200)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	PLD4 {ECO:0000303 PubMed:30111894, ECO:0000312 HGNC:HGNC:23792}
Function	5'->3' exonuclease that hydrolyzes the phosphodiester bond of single-stranded DNA (ssDNA) and RNA molecules to form nucleoside 3'- monophosphates and 5'-end 5'-hydroxy deoxyribonucleotide/ribonucleotide fragments (PubMed: <u>30111894</u> , PubMed: <u>34620855</u> , PubMed: <u>38537643</u> , PubMed: <u>39423811</u>). Partially redundant with PLD3, can cleave all four nucleotides displaying higher efficiency for ssDNA and RNA fragments initiated with uridine and guanosine residues and lower efficiency for cytidine-initiated substrates (PubMed: <u>30111894</u> , PubMed: <u>34620855</u> , PubMed: <u>38537643</u> , PubMed: <u>39423811</u>). As a result, it does not always degrade polynucleotides to the single nucleotide level, it can stall at specific sites sparing certain fragments from exonucleolytic degradation (PubMed:30111894, PubMed: <u>34620855</u> , PubMed: <u>38537643</u>)

	PubMed: <u>39423811</u>). Processes self and pathogenic ssDNA and RNA molecules that reach the endolysosomal compartment via phagocytosis or autophagy and may serve as 'danger' signals for recognition by innate immune receptors such as toll-like receptors (TLRs) (PubMed: <u>38697119</u>). Degrades mitochondrial CpG-rich ssDNA fragments to prevent TLR9 activation and autoinflammatory response, but it can cleave viral RNA to generate ligands for TLR7 activation and initiate antiviral immune responses (PubMed: <u>38697119</u>). In plasmacytoid dendritic cells, it cooperates with endonuclease RNASET2 to release 2',3'-cyclic guanosine monophosphate (2',3'-cGMP), a potent stimulatory ligand for TLR7 (PubMed: <u>38697119</u>). Produces 2',3'-cGMPs and cytidine-rich RNA fragments that occupy TLR7 ligand-binding pockets and trigger a signaling- competent state (PubMed: <u>38697119</u>). Can exert polynucleotide phosphatase activity toward 5'-phosphorylated ssDNA substrates although at a slow rate (PubMed: <u>38537643</u>). Transphosphatidylase that catalyzes the exchange with R to S stereo-inversion of the glycerol moiety between (S,R)-lysophosphatidylglycerol (LPG) and monoacylglycerol (MAG) substrates to yield (S,S)-bis(monoacylglycero)phosphate (BMP) (PubMed: <u>39423811</u>). Can synthesize a variety of (S,S)-BMPs representing the main phospholipid constituent of lysosomal intralumenal vesicle (ILV) membranes that bind acid hydrolases for lipid degradation (PubMed: <u>39423811</u>). Regulates the homeostasis and interorganellar communication of the endolysosomal system with an overall impact on cellular removal of dysfunctional organelles via autophagy as well as proper protein and lipid turnover. May play a role in myotube formation in response to ER stress (By similarity).
Cellular Location	Endoplasmic reticulum membrane {ECO:0000250 UniProtKB:Q8BG07}; Single-pass type II membrane protein {ECO:0000250 UniProtKB:Q8BG07}. Golgi apparatus, trans-Golgi network membrane {ECO:0000250 UniProtKB:Q8BG07}; Single-pass type II membrane protein {ECO:0000250 UniProtKB:Q8BG07}. Nucleus {ECO:0000250 UniProtKB:Q8BG07}. Early endosome {ECO:0000250 UniProtKB:Q8BG07}. Cytoplasmic vesicle, phagosome {ECO:0000250 UniProtKB:Q8BG07}. Lysosome Note=Activation of microglia induces translocation of PLD4 from the nucleus to the phagosomes. {ECO:0000250 UniProtKB:Q8BG07}
Tissue Location	Expressed in plasmacytoid dendritic cells and monocytes (at protein level).

Background

KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human Phospholipase D4. The exact sequence is proprietary.

Images



Western blot analysis of Phospholipase D4 expression in PC12 (A), CT26 (B), Hela (C), A549 (D) whole cell lysates.



Immunohistochemical analysis of Phospholipase D4 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.