

Anti-AMPK gamma 2 Antibody

Catalog # AP60368

Product Information

Application	WB, IHC, IF/IC
Primary Accession	Q9UGJ0
Reactivity	Human, Rat, Mk
Host	Rabbit
Clonality	Polyclonal
Calculated MW	63066

Additional Information

Gene ID	51422
Other Names	5'-AMP-activated protein kinase subunit gamma-2; AMPK gamma2; AMPK subunit gamma-2; H91620p
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human AMPK gamma 2. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500) IHC~~WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500) IF/IC~~N/A
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information




Name	PRKAG2
Function	AMP/ATP-binding subunit of AMP-activated protein kinase (AMPK), an energy sensor protein kinase that plays a key role in regulating cellular energy metabolism (PubMed: 14722619 , PubMed: 24563466). In response to reduction of intracellular ATP levels, AMPK activates energy-producing pathways and inhibits energy-consuming processes: inhibits protein, carbohydrate and lipid biosynthesis, as well as cell growth and proliferation (PubMed: 14722619 , PubMed: 24563466). AMPK acts via direct phosphorylation of metabolic enzymes, and by longer-term effects via phosphorylation of transcription regulators (PubMed: 14722619 , PubMed: 24563466). Also acts as a regulator of cellular polarity by remodeling the actin cytoskeleton; probably by indirectly activating myosin (PubMed: 14722619 , PubMed: 24563466). Gamma non-catalytic subunit mediates binding to AMP, ADP and ATP, leading to activate or inhibit AMPK: AMP-binding results in allosteric activation of alpha catalytic subunit (PRKAA1

or PRKAA2) both by inducing phosphorylation and preventing dephosphorylation of catalytic subunits (PubMed:[14722619](#), PubMed:[24563466](#)). ADP also stimulates phosphorylation, without stimulating already phosphorylated catalytic subunit (PubMed:[14722619](#), PubMed:[24563466](#)). ATP promotes dephosphorylation of catalytic subunit, rendering the AMPK enzyme inactive (PubMed:[14722619](#), PubMed:[24563466](#)).


Tissue Location

Isoform B is ubiquitously expressed except in liver and thymus. The highest level is detected in heart with abundant expression in placenta and testis

Images



Immunohistochemical analysis of AMPK gamma 2 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used



as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.