

# Anti-RPS6 (pS240) Antibody

Rabbit polyclonal antibody to RPS6 (pS240)

Catalog # AP60390

## Product Information

Application	WB, IHC, IF/IC
Primary Accession	<a href="#">P62753</a>
Other Accession	<a href="#">P62754</a>
Reactivity	Human, Mouse, Rat, Rabbit, Monkey, Chicken, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	28681

## Additional Information

Gene ID	6194
Other Names	40S ribosomal protein S6; Phosphoprotein NP33
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human RPS6 (pS240). The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500) IHC~~WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500) IF/IC~~N/A
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt


## Protein Information

Name	RPS6 {ECO:0000303   PubMed:29563586, ECO:0000312   HGNC:HGNC:10429}
Function	Component of the 40S small ribosomal subunit (PubMed: <a href="#">23636399</a> , PubMed: <a href="#">8706699</a> ). Plays an important role in controlling cell growth and proliferation through the selective translation of particular classes of mRNA (PubMed: <a href="#">17220279</a> ). Part of the small subunit (SSU) processome, first precursor of the small eukaryotic ribosomal subunit. During the assembly of the SSU processome in the nucleolus, many ribosome biogenesis factors, an RNA chaperone and ribosomal proteins associate with the nascent pre-rRNA and work in concert to generate RNA folding, modifications, rearrangements and cleavage as well as targeted degradation of pre-ribosomal RNA by the RNA exosome (PubMed: <a href="#">34516797</a> ).
Cellular Location	Cytoplasm. Nucleus, nucleolus


## Background

KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human RPS6 (pS240). The exact sequence is proprietary.


## Images



Western blot analysis of RPS6 (pS240) expression in HeLa (A), HGC27 (B) whole cell lysates.



Immunohistochemical analysis of RPS6 (pS240) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of RPS6 (pS240) staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.