

Anti-VDR (pS208) Antibody

Rabbit polyclonal antibody to VDR (pS208)

Catalog # AP60416

Product Information

Application	WB, IF/IC
Primary Accession	P11473
Other Accession	P48281
Reactivity	Human, Monkey
Host	Rabbit
Clonality	Polyclonal
Calculated MW	48289

Additional Information

Gene ID	7421
Other Names	NR1I1; Vitamin D3 receptor; VDR; 1, 25-dihydroxyvitamin D3 receptor; Nuclear receptor subfamily 1 group I member 1
Target/Specificity	Recognizes endogenous levels of VDR (pS208) protein.
Dilution	WB~~WB (1/500 - 1/1000), IF/IC (1/100 - 1/500) IF/IC~~N/A
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	VDR (HGNC:12679)
Synonyms	NR1I1
Function	<p>Nuclear receptor for calcitriol, the active form of vitamin D3 which mediates the action of this vitamin on cells (PubMed:10678179, PubMed:15728261, PubMed:16913708, PubMed:28698609, PubMed:37478846). Enters the nucleus upon vitamin D3 binding where it forms heterodimers with the retinoid X receptor/RXR (PubMed:28698609). The VDR-RXR heterodimers bind to specific response elements on DNA and activate the transcription of vitamin D3-responsive target genes (PubMed:28698609). Plays a central role in calcium homeostasis (By similarity). Also functions as a receptor for the secondary bile acid lithocholic acid (LCA) and its metabolites (PubMed:12016314, PubMed:32354638).</p> <p>Nucleus {ECO:0000255 PROSITE-ProRule:PRU00407,</p>


Cellular Location

ECO:0000269 | PubMed:12145331, ECO:0000269 | PubMed:16207705, ECO:0000269 | PubMed:28698609}. Cytoplasm Note=Localizes mainly to the nucleus (PubMed:12145331, PubMed:28698609). Translocated into the nucleus via both ligand- dependent and ligand-independent pathways; ligand-independent nuclear translocation is mediated by IPO4 (PubMed:16207705)


Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human VDR (pS208). The exact sequence is proprietary.

Images



Western blot analysis of VDR (pS208) expression in HCT116 (A), HEK293T (B) whole cell lysates.



Immunofluorescent analysis of VDR (pS208) staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.