

# Anti-CD184 Antibody

Rabbit polyclonal antibody to CD184

Catalog # AP60446

## Product Information

Application	WB
Primary Accession	<a href="#">P61073</a>
Other Accession	<a href="#">P70658</a>
Reactivity	Human, Mouse, Rat, Rabbit, Monkey, Pig, Bovine, Dog, SARS
Host	Rabbit
Clonality	Polyclonal
Calculated MW	39746

## Additional Information

Gene ID	7852
Other Names	C-X-C chemokine receptor type 4; CXC-R4; CXCR-4; FB22; Fusin; HM89; LCR1; Leukocyte-derived seven transmembrane domain receptor; LESTR; Lipopolysaccharide-associated protein 3; LAP-3; LPS-associated protein 3; NPYRL; Stromal cell-derived factor 1 receptor; SDF-1 receptor; CD184
Target/Specificity	Recognizes endogenous levels of CD184 protein.
Dilution	WB~~WB (1/500 - 1/1000)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

## Protein Information

Name	CXCR4
Function	Receptor for the C-X-C chemokine CXCL12/SDF-1 that transduces a signal by increasing intracellular calcium ion levels and enhancing MAPK1/MAPK3 activation (PubMed: <a href="#">10452968</a> , PubMed: <a href="#">18799424</a> , PubMed: <a href="#">24912431</a> , PubMed: <a href="#">28978524</a> ). Involved in the AKT signaling cascade (PubMed: <a href="#">24912431</a> ). Plays a role in regulation of cell migration, e.g. during wound healing (PubMed: <a href="#">28978524</a> ). Acts as a receptor for extracellular ubiquitin; leading to enhanced intracellular calcium ions and reduced cellular cAMP levels (PubMed: <a href="#">20228059</a> ). Binds bacterial lipopolysaccharide (LPS) et mediates LPS-induced inflammatory response, including TNF secretion by monocytes (PubMed: <a href="#">11276205</a> ). Involved in hematopoiesis and in cardiac ventricular septum formation. Also plays an essential role in vascularization of the gastrointestinal tract, probably by regulating vascular branching and/or

remodeling processes in endothelial cells. Involved in cerebellar development. In the CNS, could mediate hippocampal-neuron survival (By similarity).

#### Cellular Location

Cell membrane; Multi-pass membrane protein. Cell junction. Early endosome. Late endosome. Lysosome. Note=In unstimulated cells, diffuse pattern on plasma membrane. On agonist stimulation, colocalizes with ITCH at the plasma membrane where it becomes ubiquitinated. In the presence of antigen, distributes to the immunological synapse forming at the T- cell-APC contact area, where it localizes at the peripheral and distal supramolecular activation cluster (SMAC)

#### Tissue Location

Expressed in numerous tissues, such as peripheral blood leukocytes, spleen, thymus, spinal cord, heart, placenta, lung, liver, skeletal muscle, kidney, pancreas, cerebellum, cerebral cortex and medulla (in microglia as well as in astrocytes), brain microvascular, coronary artery and umbilical cord endothelial cells Isoform 1 is predominant in all tissues tested


## Background

---

KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human CD184. The exact sequence is proprietary.

## Images

---



Western blot analysis of CD184 expression in Hela (A), mouse lung (B), mouse spleen (C), mouse heart (D), rat spleen (E), rat heart (F) whole cell lysates.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.