

Anti-Separase Antibody

Rabbit polyclonal antibody to Separase
Catalog # AP60459

Product Information

Application	WB, IF/IC, IHC
Primary Accession	Q14674
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	233175

Additional Information

Gene ID	9700
Other Names	ESP1; KIAA0165; Separin; Caspase-like protein ESPL1; Extra spindle poles-like 1 protein; Separase
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Separase. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500) IF/IC~~N/A IHC~~WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt


Protein Information

Name	ESPL1
Synonyms	ESP1, KIAA0165
Function	Caspase-like protease, which plays a central role in the chromosome segregation by cleaving the SCC1/RAD21 subunit of the cohesin complex at the onset of anaphase. During most of the cell cycle, it is inactivated by different mechanisms.
Cellular Location	Cytoplasm. Nucleus.


Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Separase. The exact sequence is proprietary.


Images



Western blot analysis of Separase expression in HeLa (A), SGC7901 (B), HEK293T (C), CT26 (D) whole cell lysates.



Immunohistochemical analysis of Separase staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of Separase staining in HuvEc cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.